

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
21 February 2002 (21.02.2002)

PCT

(10) International Publication Number
WO 02/14483 A2

(51) International Patent Classification: C12N 9/00

(21) International Application Number: PCT/US01/25457

(22) International Filing Date: 14 August 2001 (14.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/225,493 15 August 2000 (15.08.2000) US(71) Applicant: ZYMOGENETICS INC [US/US]; 1201
Eastlake Avenue East, Seattle, WA 98102 (US).(72) Inventors: CONKLIN, Darrell, C.; Apt.#7, 2332 Minor
Avenue E, Seattle, WA 98102 (US). GAO, Zeren, 9502
179th Place NE, #3, Redmond, WA 98052 (US).(74) Agent: JONES, Phillip B, C; ZymoGenetics, Inc., 1201
Eastlake Avenue East, Seattle, WA 98102 (US).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MY, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CI,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).**Published:**

- without international search report and to be republished upon receipt of that report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HUMAN ADENOSINE DEAMINASE

(57) Abstract: Adenosine deaminase is a purine catabolic enzyme, which catalyzes the deamination of adenosine and 2-deoxyadenosine to inosine and 2-deoxyinosine, respectively. A deficiency of adenosine deaminase causes severe immunodeficiency disease. Current therapies for adenosine deficiency include bone marrow transplantation and enzyme replacement therapy. Zmim19 is a new human adenosine deaminase, which has uses in therapy, diagnostics, and industry.



WO 02/14483 A2

HUMAN ADENOSINE DEAMINASE

5

TECHNICAL FIELD

The present invention relates generally to a new gene that encodes an enzyme. In particular, the present invention relates to a novel adenosine deaminase, designated "Zmim19," and to nucleic acid molecules encoding Zmim19.

BACKGROUND OF THE INVENTION

Adenosine deaminase is a purine catabolic enzyme, which catalyzes the deamination of adenosine and 2'-deoxyadenosine to inosine and 2'-deoxyinosine, respectively. Toxic metabolites accumulate in tissues if this enzyme activity is absent.

A deficiency of adenosine deaminase causes the most severe of the immunodeficiencies in humans. Without treatment, an affected individual will die from overwhelming opportunistic infections within the first few months of life. The lack of erythrocyte adenosine deaminase is the cause of one form of severe combined immunodeficiency (SCID) (see, for example, Hirschhorn *et al.*, *Proc. Nat'l Acad. Sci. USA*, 73:213 (1976); Hong, "Inherited Immune Deficiency," in Jameson (Ed.), *Principles of Molecular Medicine*, pages 283-291 (Humana Press, Inc. 1998); Parkman *et al.*, *Ann. Rev. Med.* 51:33 (2000)). SCID is an autosomal recessively inherited disorder, which is characterized by low levels of immunoglobulins, an absence of T lymphocytes, and markedly reduced levels of B lymphocytes and natural killer cells. Adenosine deaminase deficiency accounts for about one-half of cases of autosomal recessive SCID. Decreased adenosine deaminase activity is due to either an absence of the enzyme or the decreased enzymatic activity of mutant proteins.

Current therapies for SCID include bone marrow transplantation and enzyme replacement therapy with intramuscular injections of pegademase, which is a polyethylene glycol-conjugated adenosine deaminase. Conjugation with polyethylene glycol is designed to improve circulating levels of adenosine deaminase by reducing cellular uptake, proteolysis, and immunogenicity of the native enzyme (see, for example, Herschfield *et al.* *New Eng. J. Med.* 316:589 (1987); Delgado *et al.*, *Crit. Rev. Drug Carrier Systems* 9:249 (1992)).

An ability to detect adenosine deaminase has proven useful for diagnosis of various diseases and disorders. For example, measurement of serum adenosine deaminase shows elevation in primary liver disease, secondary hepatic neoplasia, systemic lupus erythematosus, and tuberculosis (see, for example, Ellis *et al.*, *Clin. Chim. Acta* 47:75 (1973); Piras and Gakis, *Enzyme* 14:311 (1973); Collazos *et al.*, *Chest* 114:432 (1998); Stancikova *et al.*, *Clin. Exp. Rheumatol* 16:583 (1998); Kuyucu *et al.*, *J. Trop. Pediatr.* 45:245 (1999)). As described below, adenosine deaminase also has uses in industry and research.

The discovery of a new adenosine deaminase fulfills a need in the art by providing a new composition useful in diagnosis, therapy, industry, and research.

BRIEF SUMMARY OF THE INVENTION

The present invention provides a novel adenosine deaminase, designated “Zmim19.” The present invention also provides Zmim19 variant polypeptides and Zmim19 fusion proteins, as well as nucleic acid molecules encoding such polypeptides and proteins, and methods for using these nucleic acid molecules and amino acid sequences.

DETAILED DESCRIPTION OF THE INVENTION

1. Overview

The present invention provides nucleic acid molecules that encode a new human adenosine deaminase, designated as “Zmim19.” An illustrative nucleotide sequence that encodes Zmim19 is provided by SEQ ID NO:1. The encoded polypeptide has the following amino acid sequence: MIEAEEQQPC KTDIFYSELPK VELHAHLNGS ISSHTMKKLI AQKPDLEKHD QMTVIDKGKK RTLEECFQMF QTIHQLTSSP EDILMVTKDV IKEFADDGVK YLELRSTPRR ENATGMTKKT YVESILEGIK QSKQENLDID VRYLIAVDRR GGPLVAKETV KLAEEFFLST EGTVLGLDLS GDPTVGQAKD FLEPLLEAKK AGLKLALHLS EIPNQKKETQ ILLDLLPDRI GHGTFNLNGE GGSLDLVDFV RQHRIPLELC LTSNVKSQTV PSYDQHHFGF WYSIAHPSVI CTDDKGVFAT HLSQEYQLAA ETFNLTSQSV WDLSEYSEIN YIFASDSTRSE LRKKWNHLKP RVLHI (SEQ ID NO:2). Thus, the Zmim19 gene described herein encodes a polypeptide of 355 amino acids, as shown in SEQ ID NO:2. Sequence analysis indicates that the Zmim19 polypeptide includes active

site residues at His²⁰⁸, Cys²⁶⁰, Asp²⁹³, and Asp²⁹⁴. The *Zmim19* gene is expressed in uterine tissue, indicating that *Zmim19* nucleotide sequences and anti-Zmim19 antibodies can be useful for tissue differentiation.

As detailed below, the present invention provides isolated polypeptides
5 having an amino acid sequence that is at least 70%, at least 80%, or at least 90% identical to the amino acid sequence of SEQ ID NO:2, or amino acid residues 208 to 294 of SEQ ID NO:2. Particular polypeptides specifically bind with an antibody that specifically binds with a polypeptide having the amino acid sequence of SEQ ID NO:2. Certain polypeptides also can be characterized by adenosine deaminase activity. An
10 example of a polypeptide having an amino acid sequence that is at least 80% identical to the amino acid sequence of amino acid residues 208 to 294 of SEQ ID NO:2 is the amino acid sequence of amino acid residues 207 to 293 of SEQ ID NO:6.

An illustrative polypeptide is a polypeptide that comprises the amino acid sequence of SEQ ID NO:2, or that comprises amino acid residues 208 to 294 of SEQ ID
15 NO:2. Additional exemplary polypeptides include polypeptides comprising an amino acid sequence of 15, 20, or 30 contiguous amino acids of an amino acid sequence selected from the group consisting of: amino acid residues 208 to 294 of SEQ ID NO:2, and SEQ ID NO:2.

The polypeptides described herein can further comprise an affinity tag.

The present invention further provides variant Zmim19 polypeptides,
20 which have an amino acid sequence that shares an identity with the amino acid sequence of SEQ ID NO:2, selected from the group consisting of at least 70% identity, at least 80% identity, at least 90% identity, at least 95% identity, or greater than 95% identity, and wherein any difference between the amino acid sequence of the variant polypeptide
25 and the amino acid sequence of SEQ ID NO:2 is due to one or more conservative amino acid substitutions. Suitable conservative amino acid substitutions can be identified by comparison with the murine Zmim19 sequence (SEQ ID NO:6), such as Ser⁷⁸ to Thr⁷⁸, Val⁸⁵ to Ile⁸⁵, Lys¹¹⁸ to Arg¹¹⁸, and the like.

The present invention further provides antibodies and antibody fragments
30 that specifically bind with such polypeptides. Exemplary antibodies include polyclonal antibodies, murine monoclonal antibodies, humanized antibodies derived from murine monoclonal antibodies, and human monoclonal antibodies. Illustrative antibody fragments include F(ab')₂, F(ab)₂, Fab', Fab, Fv, scFv, and minimal recognition units. The present invention further includes compositions comprising a carrier and a protein,
35 peptide, polypeptide, antibody, or anti-idiotypic antibody described herein.

The present invention also provides isolated nucleic acid molecules that encode a Zmim19 polypeptide, wherein the nucleic acid molecule is selected from the

group consisting of: a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:3; a nucleic acid molecule encoding the amino acid sequence of SEQ ID NO:2; and a nucleic acid molecule that remains hybridized following stringent wash conditions to a nucleic acid molecule consisting of a nucleotide sequence selected from the group
5 consisting of: (a) the nucleotide sequence of SEQ ID NO:1, (b) nucleotides 172 to 1236 of SEQ ID NO:1, and (c) a nucleotide sequence that is the complement of the nucleotide sequence of (a) or (b).

Illustrative nucleic acid molecules include those in which any difference between the amino acid sequence encoded by the nucleic acid molecule and the
10 corresponding amino acid sequence of SEQ ID NO:2 is due to a conservative amino acid substitution. The present invention further contemplates isolated nucleic acid molecules that comprise the nucleotide sequence of SEQ ID NO:1, or nucleotides 172 to 1236 of SEQ ID NO:1.

The present invention also includes vectors and expression vectors
15 comprising such nucleic acid molecules. Such expression vectors may comprise a transcription promoter, and a transcription terminator, wherein the promoter is operably linked with the nucleic acid molecule, and wherein the nucleic acid molecule is operably linked with the transcription terminator. The present invention further includes recombinant host cells, and recombinant viruses, comprising these vectors and
20 expression vectors. Illustrative host cells include avian, bacterial, yeast, fungal, insect, mammalian, and plant cells. Recombinant host cells comprising such expression vectors can be used to produce *Zmim19* polypeptides by culturing such recombinant host cells that comprise the expression vector and that produce the *Zmim19* protein, and, optionally, isolating the *Zmim19* protein from the cultured recombinant host cells. The
25 present invention also includes the protein products of these processes.

The present invention also contemplates methods for detecting the presence of *Zmim19* RNA in a biological sample, comprising the steps of (a) contacting a *Zmim19* nucleic acid probe under hybridizing conditions with either (i) test RNA molecules isolated from the biological sample, or (ii) nucleic acid molecules synthesized
30 from the isolated RNA molecules, wherein the probe has a nucleotide sequence comprising a portion of the nucleotide sequence of SEQ ID NO:1, or its complement, and (b) detecting the formation of hybrids of the nucleic acid probe and either the test RNA molecules or the synthesized nucleic acid molecules, wherein the presence of the hybrids indicates the presence of *Zmim19* RNA in the biological sample. An example of
35 a biological sample is a human biological sample, such as a biopsy or autopsy specimen.

The present invention further provides methods for detecting the presence of *Zmim19* polypeptide in a biological sample, comprising the steps of: (a) contacting

the biological sample with an antibody or an antibody fragment that specifically binds with a polypeptide comprising the amino acid sequence of SEQ ID NO:2, wherein the contacting is performed under conditions that allow the binding of the antibody or antibody fragment to the biological sample, and (b) detecting any of the bound antibody or bound antibody fragment. Such an antibody or antibody fragment may further comprise a detectable label selected from the group consisting of radioisotope, fluorescent label, chemiluminescent label, enzyme label, bioluminescent label, and colloidal gold. An exemplary biological sample is a human biological sample.

The present invention also provides kits for performing these detection methods. For example, a kit for detection of *Zmim19* gene expression may comprise a container that comprises a nucleic acid molecule, wherein the nucleic acid molecule is selected from the group consisting of (a) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, (b) a nucleic acid molecule comprising the complement of the nucleotide sequence of SEQ ID NO:1, (c) a nucleic acid molecule that is a fragment of (a) consisting of at least eight nucleotides, and (d) a nucleic acid molecule that is a fragment of (b) consisting of at least eight nucleotides. Illustrative nucleic acid molecules include nucleic acid molecules comprising nucleotides 172 to 1236 of SEQ ID NO:1, or the complement thereof. Such a kit may also comprise a second container that comprises one or more reagents capable of indicating the presence of the nucleic acid molecule. On the other hand, a kit for detection of *Zmim19* protein may comprise a container that comprises an antibody, or an antibody fragment, that specifically binds with a polypeptide having the amino acid sequence of SEQ ID NO:2.

The present invention further provides fusion proteins comprising a *Zmim19* moiety. Examples of such fusion proteins include polypeptides comprising a *Zmim19* moiety and a cell recognition moiety. Suitable cell recognition moieties include receptor ligands, antibodies, and antibody fragments. Other types of fusion proteins include a *Zmim19* moiety and an immunoglobulin heavy chain constant region, such as a human F_C fragment. The present invention further includes isolated nucleic acid molecules that encode such fusion proteins.

The present invention also includes murine *Zmim19* polypeptides, such as the amino acid sequence of SEQ ID NO:6, and nucleic acid molecules encoding murine polypeptides, such as the nucleotide sequence of SEQ ID NO:5. Illustrative polypeptides include polypeptides that comprise the amino acid sequence of SEQ ID NO:6, or that comprise amino acid residues 207 to 293 of SEQ ID NO:6. Additional exemplary polypeptides include polypeptides comprising an amino acid sequence of 15, 20, or 30 contiguous amino acids of an amino acid sequence selected from the group consisting of: amino acid residues 207 to 293 of SEQ ID NO:6, and SEQ ID NO:6.

These and other aspects of the invention will become evident upon reference to the following detailed description. In addition, various references are identified below.

2. Definitions

In the description that follows, a number of terms are used extensively. The following definitions are provided to facilitate understanding of the invention.

As used herein, "nucleic acid" or "nucleic acid molecule" refers to polynucleotides, such as deoxyribonucleic acid (DNA) or ribonucleic acid (RNA), oligonucleotides, fragments generated by the polymerase chain reaction (PCR), and fragments generated by any of ligation, scission, endonuclease action, and exonuclease action. Nucleic acid molecules can be composed of monomers that are naturally-occurring nucleotides (such as DNA and RNA), or analogs of naturally-occurring nucleotides (e.g., α -enantiomeric forms of naturally-occurring nucleotides), or a combination of both. Modified nucleotides can have alterations in sugar moieties and/or in pyrimidine or purine base moieties. Sugar modifications include, for example, replacement of one or more hydroxyl groups with halogens, alkyl groups, amines, and azido groups, or sugars can be functionalized as ethers or esters. Moreover, the entire sugar moiety can be replaced with sterically and electronically similar structures, such as aza-sugars and carbocyclic sugar analogs. Examples of modifications in a base moiety include alkylated purines and pyrimidines, acylated purines or pyrimidines, or other well-known heterocyclic substitutes. Nucleic acid monomers can be linked by phosphodiester bonds or analogs of such linkages. Analogs of phosphodiester linkages include phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoranilidate, phosphoramidate, and the like. The term "nucleic acid molecule" also includes so-called "peptide nucleic acids," which comprise naturally-occurring or modified nucleic acid bases attached to a polyamide backbone. Nucleic acids can be either single stranded or double stranded.

The term "complement of a nucleic acid molecule" refers to a nucleic acid molecule having a complementary nucleotide sequence and reverse orientation as compared to a reference nucleotide sequence. For example, the sequence 5' ATGCACGGG 3' is complementary to 5' CCCGTGCAT 3'.

The term "contig" denotes a nucleic acid molecule that has a contiguous stretch of identical or complementary sequence to another nucleic acid molecule.

Contiguous sequences are said to "overlap" a given stretch of a nucleic acid molecule either in their entirety or along a partial stretch of the nucleic acid molecule.

The term "degenerate nucleotide sequence" denotes a sequence of nucleotides that includes one or more degenerate codons as compared to a reference
5 nucleic acid molecule that encodes a polypeptide. Degenerate codons contain different triplets of nucleotides, but encode the same amino acid residue (*i.e.*, GAU and GAC triplets each encode Asp).

The term "structural gene" refers to a nucleic acid molecule that is transcribed into messenger RNA (mRNA), which is then translated into a sequence of
10 amino acids characteristic of a specific polypeptide.

An "isolated nucleic acid molecule" is a nucleic acid molecule that is not integrated in the genomic DNA of an organism. For example, a DNA molecule that encodes a growth factor that has been separated from the genomic DNA of a cell is an isolated DNA molecule. Another example of an isolated nucleic acid molecule is a
15 chemically-synthesized nucleic acid molecule that is not integrated in the genome of an organism. A nucleic acid molecule that has been isolated from a particular species is smaller than the complete DNA molecule of a chromosome from that species.

A "nucleic acid molecule construct" is a nucleic acid molecule, either single- or double-stranded, that has been modified through human intervention to contain
20 segments of nucleic acid combined and juxtaposed in an arrangement not existing in nature.

"Linear DNA" denotes non-circular DNA molecules having free 5' and 3' ends. Linear DNA can be prepared from closed circular DNA molecules, such as plasmids, by enzymatic digestion or physical disruption.

"Complementary DNA (cDNA)" is a single-stranded DNA molecule that is
25 formed from an mRNA template by the enzyme reverse transcriptase. Typically, a primer complementary to portions of mRNA is employed for the initiation of reverse transcription. Those skilled in the art also use the term "cDNA" to refer to a double-stranded DNA molecule consisting of such a single-stranded DNA molecule and its complementary DNA
30 strand. The term "cDNA" also refers to a clone of a cDNA molecule synthesized from an RNA template.

A "promoter" is a nucleotide sequence that directs the transcription of a structural gene. Typically, a promoter is located in the 5' non-coding region of a gene, proximal to the transcriptional start site of a structural gene. Sequence elements within
35 promoters that function in the initiation of transcription are often characterized by consensus nucleotide sequences. These promoter elements include RNA polymerase binding sites, TATA sequences, CAAT sequences, differentiation-specific elements

(DSEs; McGehee *et al.*, *Mol. Endocrinol.* 7:551 (1993)), cyclic AMP response elements (CREs), serum response elements (SREs; Treisman, *Seminars in Cancer Biol.* 1:47 (1990)), glucocorticoid response elements (GREs), and binding sites for other transcription factors, such as CRE/ATF (O'Reilly *et al.*, *J. Biol. Chem.* 267:19938 (1992)), AP2 (Ye *et al.*, *J. Biol. Chem.* 269:25728 (1994)), SP1, cAMP response element binding protein (CREB; Loeken, *Gene Expr.* 3:253 (1993)) and octamer factors (see, in general, Watson *et al.*, eds., *Molecular Biology of the Gene*, 4th ed. (The Benjamin/Cummings Publishing Company, Inc. 1987), and Lemaigre and Rousseau, *Biochem. J.* 303:1 (1994)). If a promoter is an inducible promoter, then the rate of transcription increases in response to an inducing agent. In contrast, the rate of transcription is not regulated by an inducing agent if the promoter is a constitutive promoter. Repressible promoters are also known.

A "core promoter" contains essential nucleotide sequences for promoter function, including the TATA box and start of transcription. By this definition, a core promoter may or may not have detectable activity in the absence of specific sequences that may enhance the activity or confer tissue specific activity.

A "regulatory element" is a nucleotide sequence that modulates the activity of a core promoter. For example, a regulatory element may contain a nucleotide sequence that binds with cellular factors enabling transcription exclusively or preferentially in particular cells, tissues, or organelles. These types of regulatory elements are normally associated with genes that are expressed in a "cell-specific," "tissue-specific," or "organelle-specific" manner.

An "enhancer" is a type of regulatory element that can increase the efficiency of transcription, regardless of the distance or orientation of the enhancer relative to the start site of transcription.

"Heterologous DNA" refers to a DNA molecule, or a population of DNA molecules, that does not exist naturally within a given host cell. DNA molecules heterologous to a particular host cell may contain DNA derived from the host cell species (*i.e.*, endogenous DNA) so long as that host DNA is combined with non-host DNA (*i.e.*, exogenous DNA). For example, a DNA molecule containing a non-host DNA segment encoding a polypeptide operably linked to a host DNA segment comprising a transcription promoter is considered to be a heterologous DNA molecule. Conversely, a heterologous DNA molecule can comprise an endogenous gene operably linked with an exogenous promoter. As another illustration, a DNA molecule comprising a gene derived from a wild-type cell is considered to be heterologous DNA if that DNA molecule is introduced into a mutant cell that lacks the wild-type gene.

A "polypeptide" is a polymer of amino acid residues joined by peptide bonds, whether produced naturally or synthetically. Polypeptides of less than about 10 amino acid residues are commonly referred to as "peptides."

5 A "protein" is a macromolecule comprising one or more polypeptide chains. A protein may also comprise non-peptidic components, such as carbohydrate groups. Carbohydrates and other non-peptidic substituents may be added to a protein by the cell in which the protein is produced, and will vary with the type of cell. Proteins are defined herein in terms of their amino acid backbone structures; substituents such as carbohydrate groups are generally not specified, but may be present nonetheless.

10 A peptide or polypeptide encoded by a non-host DNA molecule is a "heterologous" peptide or polypeptide.

An "integrated genetic element" is a segment of DNA that has been incorporated into a chromosome of a host cell after that element is introduced into the cell through human manipulation. Within the present invention, integrated genetic elements are most commonly derived from linearized plasmids that are introduced into the cells by electroporation or other techniques. Integrated genetic elements are passed from the original host cell to its progeny.

15 A "cloning vector" is a nucleic acid molecule, such as a plasmid, cosmid, or bacteriophage, which has the capability of replicating autonomously in a host cell. Cloning vectors typically contain one or a small number of restriction endonuclease recognition sites that allow insertion of a nucleic acid molecule in a determinable fashion without loss of an essential biological function of the vector, as well as nucleotide sequences encoding a marker gene that is suitable for use in the identification and selection of cells transformed with the cloning vector. Marker genes typically include genes that provide tetracycline resistance or ampicillin resistance.

25 An "expression vector" is a nucleic acid molecule encoding a gene that is expressed in a host cell. Typically, an expression vector comprises a transcription promoter, a gene, and a transcription terminator. Gene expression is usually placed under the control of a promoter, and such a gene is said to be "operably linked to" the promoter. Similarly, a regulatory element and a core promoter are operably linked if the regulatory element modulates the activity of the core promoter.

30 A "recombinant host" is a cell that contains a heterologous nucleic acid molecule, such as a cloning vector or expression vector. In the present context, an example of a recombinant host is a cell that produces Zmim19 from an expression vector. In contrast, Zmim19 can be produced by a cell that is a "natural source" of Zmim19 (*e.g.* uterine tissue), and that lacks an expression vector.

"Integrative transformants" are recombinant host cells, in which heterologous DNA has become integrated into the genomic DNA of the cells.

A "fusion protein" is a hybrid protein expressed by a nucleic acid molecule comprising nucleotide sequences of at least two genes. For example, a fusion protein can comprise at least part of a Zmim19 polypeptide fused with a polypeptide that binds an affinity matrix. Such a fusion protein provides a means to isolate large quantities of Zmim19 using affinity chromatography.

The term "receptor" denotes a cell-associated protein that binds to a bioactive molecule termed a "ligand." This interaction mediates the effect of the ligand on the cell. Receptors can be membrane bound, cytosolic or nuclear; monomeric (*e.g.*, thyroid stimulating hormone receptor, beta-adrenergic receptor) or multimeric (*e.g.*, PDGF receptor, growth hormone receptor, IL-3 receptor, GM-CSF receptor, G-CSF receptor, erythropoietin receptor and IL-6 receptor). Membrane-bound receptors are characterized by a multi-domain structure comprising an extracellular ligand-binding domain and an intracellular effector domain that is typically involved in signal transduction. In certain membrane-bound receptors, the extracellular ligand-binding domain and the intracellular effector domain are located in separate polypeptides that comprise the complete functional receptor.

In general, the binding of ligand to receptor results in a conformational change in the receptor that causes an interaction between the effector domain and other molecule(s) in the cell, which in turn leads to an alteration in the metabolism of the cell. Metabolic events that are often linked to receptor-ligand interactions include gene transcription, phosphorylation, dephosphorylation, increases in cyclic AMP production, mobilization of cellular calcium, mobilization of membrane lipids, cell adhesion, hydrolysis of inositol lipids and hydrolysis of phospholipids.

The term "secretory signal sequence" denotes a nucleotide sequence that encodes a peptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger polypeptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

An "isolated polypeptide" is a polypeptide that is essentially free from contaminating cellular components, such as carbohydrate, lipid, or other proteinaceous impurities associated with the polypeptide in nature. Typically, a preparation of isolated polypeptide contains the polypeptide in a highly purified form, *i.e.*, at least about 80% pure, at least about 90% pure, at least about 95% pure, greater than 95% pure, or greater than 99% pure. One way to show that a particular protein preparation contains an isolated polypeptide is by the appearance of a single band following sodium dodecyl

sulfate (SDS)-polyacrylamide gel electrophoresis of the protein preparation and Coomassie Brilliant Blue staining of the gel. However, the term "isolated" does not exclude the presence of the same polypeptide in alternative physical forms, such as dimers or alternatively glycosylated or derivatized forms.

5 The terms "amino-terminal" and "carboxyl-terminal" are used herein to denote positions within polypeptides. Where the context allows, these terms are used with reference to a particular sequence or portion of a polypeptide to denote proximity or relative position. For example, a certain sequence positioned carboxyl-terminal to a reference sequence within a polypeptide is located proximal to the carboxyl terminus of
10 the reference sequence, but is not necessarily at the carboxyl terminus of the complete polypeptide.

The term "expression" refers to the biosynthesis of a gene product. For example, in the case of a structural gene, expression involves transcription of the structural gene into mRNA and the translation of mRNA into one or more polypeptides.

15 The term "splice variant" is used herein to denote alternative forms of RNA transcribed from a gene. Splice variation arises naturally through use of alternative splicing sites within a transcribed RNA molecule, or less commonly between separately transcribed RNA molecules, and may result in several mRNAs transcribed from the same gene. Splice variants may encode polypeptides having altered amino acid
20 sequence. The term splice variant is also used herein to denote a polypeptide encoded by a splice variant of an mRNA transcribed from a gene.

As used herein, the term "immunomodulator" includes cytokines, stem cell growth factors, lymphotoxins, co-stimulatory molecules, hematopoietic factors, and synthetic analogs of these molecules.

25 The term "complement/anti-complement pair" denotes non-identical moieties that form a non-covalently associated, stable pair under appropriate conditions. For instance, biotin and avidin (or streptavidin) are prototypical members of a complement/anti-complement pair. Other exemplary complement/anti-complement pairs include receptor/ligand pairs, antibody/antigen (or hapten or epitope) pairs,
30 sense/antisense polynucleotide pairs, and the like. Where subsequent dissociation of the complement/anti-complement pair is desirable, the complement/anti-complement pair preferably has a binding affinity of less than 10^9 M^{-1} .

An "anti-idiotypic antibody" is an antibody that binds with the variable region domain of an immunoglobulin. In the present context, an anti-idiotypic antibody
35 binds with the variable region of an anti-Zmim19 antibody, and thus, an anti-idiotypic antibody mimics an epitope of Zmim19. Particular Zmim19 anti-idiotypic antibodies possess adenosine deaminase activity.

An "antibody fragment" is a portion of an antibody such as F(ab')₂, F(ab)₂, Fab', Fab, and the like. Regardless of structure, an antibody fragment binds with the same antigen that is recognized by the intact antibody. For example, an anti-Zmim19 monoclonal antibody fragment binds with an epitope of Zmim19.

- 5 The term "antibody fragment" also includes a synthetic or a genetically engineered polypeptide that binds to a specific antigen, such as polypeptides consisting of the light chain variable region, "Fv" fragments consisting of the variable regions of the heavy and light chains, recombinant single chain polypeptide molecules in which light and heavy variable regions are connected by a peptide linker ("scFv proteins"), and minimal
10 recognition units consisting of the amino acid residues that mimic the hypervariable region.

A "chimeric antibody" is a recombinant protein that contains the variable domains and complementary determining regions derived from a rodent antibody, while the remainder of the antibody molecule is derived from a human antibody.

- "Humanized antibodies" are recombinant proteins in which murine
15 complementarity determining regions of a monoclonal antibody have been transferred from heavy and light variable chains of the murine immunoglobulin into a human variable domain.

- As used herein, a "therapeutic agent" is a molecule or atom, which is conjugated to an antibody moiety to produce a conjugate which is useful for therapy.
20 Examples of therapeutic agents include drugs, toxins, immunomodulators, chelators, boron compounds, photoactive agents or dyes, and radioisotopes.

- A "detectable label" is a molecule or atom, which can be conjugated to an antibody moiety to produce a molecule useful for diagnosis. Examples of detectable labels include chelators, photoactive agents, radioisotopes, fluorescent agents,
25 paramagnetic ions, or other marker moieties.

- The term "affinity tag" is used herein to denote a polypeptide segment that can be attached to a second polypeptide to provide for purification or detection of the second polypeptide or provide sites for attachment of the second polypeptide to a substrate. In principal, any peptide or protein for which an antibody or other specific
30 binding agent is available can be used as an affinity tag. Affinity tags include a polyhistidine tract, protein A (Nilsson *et al.*, *EMBO J.* 4:1075 (1985); Nilsson *et al.*, *Methods Enzymol.* 198:3 (1991)), glutathione S transferase (Smith and Johnson, *Gene* 67:31 (1988)), Glu-Glu affinity tag (Grussenmeyer *et al.*, *Proc. Natl. Acad. Sci. USA* 82:7952 (1985)), substance P, FLAG peptide (Hopp *et al.*, *Biotechnology* 6:1204 (1988)),
35 streptavidin binding peptide, or other antigenic epitope or binding domain. See, in general, Ford *et al.*, *Protein Expression and Purification* 2:95 (1991). Nucleic acid

molecules encoding affinity tags are available from commercial suppliers (e.g., Pharmacia Biotech, Piscataway, NJ).

A “naked antibody” is an entire antibody, as opposed to an antibody fragment, which is not conjugated with a therapeutic agent. Naked antibodies include both polyclonal and monoclonal antibodies, as well as certain recombinant antibodies, such as chimeric and humanized antibodies.

As used herein, the term “antibody component” includes both an entire antibody and an antibody fragment.

An “immunoconjugate” is a conjugate of an antibody component with a Zmim19 moiety. Such immunoconjugates can be produced by chemically linking an antibody component and a Zmim19 moiety. For example, immunoconjugates can be prepared by indirectly conjugating a Zmim19 moiety to an antibody component (see, for example, Shih *et al.*, *Int. J. Cancer* 41:832 (1988); Shih *et al.*, *Int. J. Cancer* 46:1101 (1990); Shih *et al.*, U.S. patent No. 5,057,313). Immunoconjugates can also be prepared by directly conjugating an antibody component with a Zmim19 moiety. As an illustration, a Zmim19 moiety is directly attached to an oxidized antibody component, a Zmim19 moiety is attached at the hinge region of a reduced antibody component via disulfide bond formation, and the like. General techniques for such conjugation are well-known in the art. See, for example, Wong, *Chemistry Of Protein Conjugation And Cross-Linking* (CRC Press 1991), Upeslakis *et al.*, “Modification of Antibodies by Chemical Methods,” In: *Monoclonal Antibodies: Principles And Applications*, Birch *et al.* (Eds.), pages 187-230. (Wiley-Liss, Inc. 1995), Price, “Production and Characterization of Synthetic Peptide-Derived Antibodies,” in *Monoclonal Antibodies: Production, Engineering And Clinical Application*, Ritter *et al.* (Eds.), pages 60-84 (Cambridge University Press 1995).

As used herein, the term “antibody fusion protein” refers to a recombinant molecule that comprises an antibody component and a Zmim19 moiety.

An “adenosine deaminase targeting composition,” or a “Zmim19 targeting composition” comprises a Zmim19 moiety (e.g., Zmim19, a Zmim19 fragment, a molecule having Zmim19 activity, and the like) and a recognition molecule. Illustrative recognition molecules include antibodies, antibody components, receptor ligands, and other members of a complement/anti-complement pair. The association between the Zmim19 moiety and the recognition molecule can be covalent or noncovalent. For example, the association between a Zmim19 moiety and a recognition molecule in immunoconjugates and antibody fusion proteins is covalent, while liposomes can comprise a Zmim19 moiety and a recognition molecule in a noncovalent association.

A "tumor associated antigen" is a protein normally not expressed, or expressed at lower levels, by a normal counterpart cell. Examples of tumor associated antigens include α -fetoprotein, carcinoembryonic antigen, and Her-2/neu. Many other examples of tumor associated antigens are known to those of skill in the art. See, for example, Urban *et al.*, *Ann. Rev. Immunol.* 10:617 (1992), Garrett and Sell (Eds.), *Cellular Cancer Markers* (Humana Press 1995), Hanausek and Walaszek (Eds.), *Tumor Marker Protocols* (Humana Press 1998), and Jaffee, *Ann. N. Y. Acad. Sci.* 886:67 (1999).

As used herein, an "infectious agent" denotes both microbes and parasites. A "microbe" includes viruses, bacteria, rickettsia, mycoplasma, protozoa, fungi and like microorganisms. A "parasite" denotes infectious, generally microscopic or very small multicellular invertebrates, or ova or juvenile forms thereof, which are susceptible to immune-mediated clearance or lytic or phagocytic destruction, such as malarial parasites, spirochetes, and the like.

An "infectious agent antigen" is an antigen associated with an infectious agent.

A "target polypeptide" or a "target peptide" is an amino acid sequence that comprises at least one epitope, and that is expressed on a target cell, such as a tumor cell, or a cell that carries an infectious agent antigen. T cells recognize peptide epitopes presented by a major histocompatibility complex molecule to a target polypeptide or target peptide and typically lyse the target cell or recruit other immune cells to the site of the target cell, thereby killing the target cell.

An "antigenic peptide" is a peptide that will bind a major histocompatibility complex molecule to form an MHC-peptide complex which is recognized by a T cell, thereby inducing a cytotoxic lymphocyte response upon presentation to the T cell. Thus, antigenic peptides are capable of binding to an appropriate major histocompatibility complex molecule and inducing a cytotoxic T cells response, such as cell lysis or specific cytokine release against the target cell which binds or expresses the antigen. The antigenic peptide can be bound in the context of a class I or class II major histocompatibility complex molecule, on an antigen presenting cell or on a target cell.

In eukaryotes, RNA polymerase II catalyzes the transcription of a structural gene to produce mRNA. A nucleic acid molecule can be designed to contain an RNA polymerase II template in which the RNA transcript has a sequence that is complementary to that of a specific mRNA. The RNA transcript is termed an "anti-sense RNA" and a nucleic acid molecule that encodes the anti-sense RNA is termed an "anti-sense gene." Anti-sense RNA molecules are capable of binding to mRNA molecules, resulting in an inhibition of mRNA translation.

An “anti-sense oligonucleotide specific for *Znim19*” or an “*Znim19* anti-sense oligonucleotide” is an oligonucleotide having a sequence (a) capable of forming a stable triplex with a portion of the *Znim19* gene, or (b) capable of forming a stable duplex with a portion of an mRNA transcript of the *Znim19* gene.

5 A “ribozyme” is a nucleic acid molecule that contains a catalytic center. The term includes RNA enzymes, self-splicing RNAs, self-cleaving RNAs, and nucleic acid molecules that perform these catalytic functions. A nucleic acid molecule that encodes a ribozyme is termed a “ribozyme gene.”

10 An “external guide sequence” is a nucleic acid molecule that directs the endogenous ribozyme, RNase P, to a particular species of intracellular mRNA, resulting in the cleavage of the mRNA by RNase P. A nucleic acid molecule that encodes an external guide sequence is termed an “external guide sequence gene.”

The term “variant *Znim19* gene” refers to nucleic acid molecules that encode a polypeptide having an amino acid sequence that is a modification of SEQ ID NO:2. Such variants include naturally-occurring polymorphisms of *Znim19* genes, as well as synthetic genes that contain conservative amino acid substitutions of the amino acid sequence of SEQ ID NO:2. Additional variant forms of *Znim19* genes are nucleic acid molecules that contain insertions or deletions of the nucleotide sequences described herein. A variant *Znim19* gene can be identified by determining whether the gene hybridizes with a nucleic acid molecule consisting of the nucleotide sequence of SEQ ID NO:1, or its complement, under stringent conditions.

Alternatively, variant *Znim19* genes can be identified by sequence comparison. Two amino acid sequences have “100% amino acid sequence identity” if the amino acid residues of the two amino acid sequences are the same when aligned for maximal correspondence. Similarly, two nucleotide sequences have “100% nucleotide sequence identity” if the nucleotide residues of the two nucleotide sequences are the same when aligned for maximal correspondence. Sequence comparisons can be performed using standard software programs such as those included in the LASERGENE bioinformatics computing suite, which is produced by DNASTAR (Madison, Wisconsin). Other methods for comparing two nucleotide or amino acid sequences by determining optimal alignment are well-known to those of skill in the art (see, for example, Peruski and Peruski, *The Internet and the New Biology: Tools for Genomic and Molecular Research* (ASM Press, Inc. 1997), Wu *et al.* (eds.), “Information Superhighway and Computer Databases of Nucleic Acids and Proteins,” in *Methods in Gene Biotechnology*, pages 123-151 (CRC Press, Inc. 1997), and Bishop (ed.), *Guide to Human Genome Computing*, 2nd Edition (Academic Press, Inc. 1998)). Particular methods for determining sequence identity are described below.

Regardless of the particular method used to identify a variant *Zmim19* gene or variant *Zmim19* polypeptide, a variant gene or polypeptide encoded by a variant gene may be characterized by at least one of: the ability to bind specifically to an anti-*Zmim19* antibody, and adenosine deaminase activity.

5 The term "allelic variant" is used herein to denote any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in phenotypic polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequence. The term allelic variant is
10 also used herein to denote a protein encoded by an allelic variant of a gene.

The term "ortholog" denotes a polypeptide or protein obtained from one species that is the functional counterpart of a polypeptide or protein from a different species. Sequence differences among orthologs are the result of speciation.

"Paralogs" are distinct but structurally related proteins made by an
15 organism. Paralogs are believed to arise through gene duplication. For example, α -globin, β -globin, and myoglobin are paralogs of each other.

The present invention includes functional fragments of *Zmim19* genes. Within the context of this invention, a "functional fragment" of a *Zmim19* gene refers to a nucleic acid molecule that encodes a portion of a *Zmim19* polypeptide which
20 specifically binds with an anti-*Zmim19* antibody or possesses adenosine deaminase activity. For example, a functional fragment of a *Zmim19* gene described herein comprises a portion of the nucleotide sequence of SEQ ID NO:1, and encodes a polypeptide that specifically binds with an anti-*Zmim19* antibody.

Due to the imprecision of standard analytical methods, molecular weights
25 and lengths of polymers are understood to be approximate values. When such a value is expressed as "about" X or "approximately" X, the stated value of X will be understood to be accurate to $\pm 10\%$.

3. *Production of Nucleic Acid Molecules Encoding Zmim19*

30 Nucleic acid molecules encoding human *Zmim19* can be obtained by screening a human cDNA or genomic library using polynucleotide probes based upon SEQ ID NO:1. These techniques are standard and well-established.

As an illustration, a nucleic acid molecule that encodes a human *Zmim19* gene can be isolated from a human cDNA library. In this case, the first step would be to
35 prepare the cDNA library by isolating RNA from tissue (*e.g.*, uterine tissue), using methods well-known to those of skill in the art. In general, RNA isolation techniques must

provide a method for breaking cells, a means of inhibiting RNase-directed degradation of RNA, and a method of separating RNA from DNA, protein, and polysaccharide contaminants. For example, total RNA can be isolated by freezing tissue in liquid nitrogen, grinding the frozen tissue with a mortar and pestle to lyse the cells, extracting the ground
5 tissue with a solution of phenol/chloroform to remove proteins, and separating RNA from the remaining impurities by selective precipitation with lithium chloride (see, for example, Ausubel *et al.* (eds.), *Short Protocols in Molecular Biology*, 3rd Edition, pages 4-1 to 4-6 (John Wiley & Sons 1995) ["Ausubel (1995)"]; Wu *et al.*, *Methods in Gene Biotechnology*, pages 33-41 (CRC Press, Inc. 1997) ["Wu (1997)"].

10 Alternatively, total RNA can be isolated from tissue by extracting ground tissue with guanidinium isothiocyanate, extracting with organic solvents, and separating RNA from contaminants using differential centrifugation (see, for example, Chirgwin *et al.*, *Biochemistry* 18:52 (1979); Ausubel (1995) at pages 4-1 to 4-6; Wu (1997) at pages 33-41).

15 In order to construct a cDNA library, poly(A)⁺ RNA must be isolated from a total RNA preparation. Poly(A)⁺ RNA can be isolated from total RNA using the standard technique of oligo(dT)-cellulose chromatography (see, for example, Aviv and Leder, *Proc. Nat'l Acad. Sci. USA* 69:1408 (1972); Ausubel (1995) at pages 4-11 to 4-12).

20 Double-stranded cDNA molecules are synthesized from poly(A)⁺ RNA using techniques well-known to those in the art. (see, for example, Wu (1997) at pages 41-46). Moreover, commercially available kits can be used to synthesize double-stranded cDNA molecules. For example, such kits are available from Life Technologies, Inc. (Gaithersburg, MD), CLONTECH Laboratories, Inc. (Palo Alto, CA), Promega Corporation (Madison, WI) and STRATAGENE (La Jolla, CA).

25 Various cloning vectors are appropriate for the construction of a cDNA library. For example, a cDNA library can be prepared in a vector derived from bacteriophage, such as a λ gt10 vector. See, for example, Huynh *et al.*, "Constructing and Screening cDNA Libraries in λ gt10 and λ gt11," in *DNA Cloning: A Practical Approach Vol. I*, Glover (ed.), page 49 (IRL Press, 1985); Wu (1997) at pages 47-52.

30 Alternatively, double-stranded cDNA molecules can be inserted into a plasmid vector, such as a PBLUESCRIPT vector (STRATAGENE; La Jolla, CA), a LAMDA GEM-4 (Promega Corp.) or other commercially available vectors. Suitable cloning vectors also can be obtained from the American Type Culture Collection (Manassas, VA).

35 To amplify the cloned cDNA molecules, the cDNA library is inserted into a prokaryotic host, using standard techniques. For example, a cDNA library can be

introduced into competent *E. coli* DH5 cells, which can be obtained, for example, from Life Technologies, Inc. (Gaithersburg, MD).

A human genomic library can be prepared by means well-known in the art (see, for example, Ausubel (1995) at pages 5-1 to 5-6; Wu (1997) at pages 307-327).

- 5 Genomic DNA can be isolated by lysing tissue with the detergent Sarkosyl, digesting the lysate with proteinase K, clearing insoluble debris from the lysate by centrifugation, precipitating nucleic acid from the lysate using isopropanol, and purifying resuspended DNA on a cesium chloride density gradient.

- DNA fragments that are suitable for the production of a genomic library can
10 be obtained by the random shearing of genomic DNA or by the partial digestion of genomic DNA with restriction endonucleases. Genomic DNA fragments can be inserted into a vector, such as a bacteriophage or cosmid vector, in accordance with conventional techniques, such as the use of restriction enzyme digestion to provide appropriate termini, the use of alkaline phosphatase treatment to avoid undesirable joining of DNA molecules,
15 and ligation with appropriate ligases. Techniques for such manipulation are well-known in the art (see, for example, Ausubel (1995) at pages 5-1 to 5-6; Wu (1997) at pages 307-327).

- Nucleic acid molecules that encode a human *Zmim19* gene can also be obtained using the polymerase chain reaction (PCR) with oligonucleotide primers having nucleotide sequences that are based upon the nucleotide sequences of the human *Zmim19*
20 gene, as described herein. General methods for screening libraries with PCR are provided by, for example, Yu *et al.*, "Use of the Polymerase Chain Reaction to Screen Phage Libraries," in *Methods in Molecular Biology, Vol. 15: PCR Protocols: Current Methods and Applications*, White (ed.), pages 211-215 (Humana Press, Inc. 1993). Moreover, techniques for using PCR to isolate related genes are described by, for
25 example, Preston, "Use of Degenerate Oligonucleotide Primers and the Polymerase Chain Reaction to Clone Gene Family Members," in *Methods in Molecular Biology, Vol. 15: PCR Protocols: Current Methods and Applications*, White (ed.), pages 317-337 (Humana Press, Inc. 1993).

- Alternatively, human genomic libraries can be obtained from commercial
30 sources such as Research Genetics (Huntsville, AL) and the American Type Culture Collection (Manassas, VA).

A library containing cDNA or genomic clones can be screened with one or more polynucleotide probes based upon SEQ ID NO:1, using standard methods (see, for example, Ausubel (1995) at pages 6-1 to 6-11).

- 35 Anti-*Zmim19* antibodies, produced as described below, can also be used to isolate DNA sequences that encode human *Zmim19* genes from cDNA libraries. For example, the antibodies can be used to screen λ gt11 expression libraries, or the

antibodies can be used for immunoscreening following hybrid selection and translation (see, for example, Ausubel (1995) at pages 6-12 to 6-16; Margolis *et al.*, "Screening λ expression libraries with antibody and protein probes," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover *et al.* (eds.), pages 1-14 (Oxford University Press 1995)).

5 As an alternative, a *Zmim19* gene can be obtained by synthesizing nucleic acid molecules using mutually priming long oligonucleotides and the nucleotide sequences described herein (see, for example, Ausubel (1995) at pages 8-8 to 8-9). Established techniques using the polymerase chain reaction provide the ability to synthesize DNA molecules at least two kilobases in length (Adang *et al.*, *Plant Molec. Biol.* 21:1131 (1993), Bambot *et al.*, *PCR Methods and Applications* 2:266 (1993),
10 Dillon *et al.*, "Use of the Polymerase Chain Reaction for the Rapid Construction of Synthetic Genes," in *Methods in Molecular Biology, Vol. 15: PCR Protocols: Current Methods and Applications*, White (ed.), pages 263-268, (Humana Press, Inc. 1993), and Holowachuk *et al.*, *PCR Methods Appl.* 4:299 (1995)).

15 The nucleic acid molecules of the present invention can also be synthesized with "gene machines" using protocols such as the phosphoramidite method. If chemically-synthesized double stranded DNA is required for an application such as the synthesis of a gene or a gene fragment, then each complementary strand is made separately. The production of short genes (60 to 80 base pairs) is technically
20 straightforward and can be accomplished by synthesizing the complementary strands and then annealing them. For the production of longer genes (>300 base pairs), however, special strategies may be required, because the coupling efficiency of each cycle during chemical DNA synthesis is seldom 100%. To overcome this problem, synthetic genes (double-stranded) are assembled in modular form from single-stranded fragments that
25 are from 20 to 100 nucleotides in length. For reviews on polynucleotide synthesis, see, for example, Glick and Pasternak, *Molecular Biotechnology, Principles and Applications of Recombinant DNA* (ASM Press 1994), Itakura *et al.*, *Annu. Rev. Biochem.* 53:323 (1984), and Climie *et al.*, *Proc. Nat'l Acad. Sci. USA* 87:633 (1990).

The sequence of a *Zmim19* cDNA or *Zmim19* genomic fragment can be
30 determined using standard methods. *Zmim19* polynucleotide sequences disclosed herein can also be used as probes or primers to clone 5' non-coding regions of a *Zmim19* gene. Promoter elements from a *Zmim19* gene can be used to direct the expression of heterologous genes in, for example, uterine tissue of transgenic animals or patients undergoing gene therapy. The identification of genomic fragments containing a *Zmim19*
35 promoter or regulatory element can be achieved using well-established techniques, such as deletion analysis (see, generally, Ausubel (1995)).

Cloning of 5' flanking sequences also facilitates production of *Zmim19* proteins by "gene activation," as disclosed in U.S. Patent No. 5,641,670. Briefly, expression of an endogenous *Zmim19* gene in a cell is altered by introducing into the *Zmim19* locus a DNA construct comprising at least a targeting sequence, a regulatory
5 sequence, an exon, and an unpaired splice donor site. The targeting sequence is a *Zmim19* 5' non-coding sequence that permits homologous recombination of the construct with the endogenous *Zmim19* locus, whereby the sequences within the construct become operably linked with the endogenous *Zmim19* coding sequence. In this way, an endogenous *Zmim19* promoter can be replaced or supplemented with other regulatory
10 sequences to provide enhanced, tissue-specific, or otherwise regulated expression.

4. *Production of Zmim19 Variants*

The present invention provides a variety of nucleic acid molecules, including DNA and RNA molecules that encode the *Zmim19* polypeptides disclosed
15 herein. Those skilled in the art will readily recognize that, in view of the degeneracy of the genetic code, considerable sequence variation is possible among these polynucleotide molecules. SEQ ID NO:3 is a degenerate nucleotide sequence that encompasses all nucleic acid molecules that encode the *Zmim19* polypeptide of SEQ ID NO:2. Those
20 skilled in the art will recognize that the degenerate sequence of SEQ ID NO:3 also provides all RNA sequences encoding SEQ ID NO:2, by substituting U for T. Thus, the present invention contemplates *Zmim19* polypeptide-encoding nucleic acid molecules comprising nucleotides 172 to 1236 of SEQ ID NO:1, and their RNA equivalents. The
present invention also includes murine *Zmim19* polypeptide-encoding nucleic acid molecules comprising nucleotides 366 to 1445 of SEQ ID NO:5, and their RNA
25 equivalents.

Table 1 sets forth the one-letter codes used within SEQ ID NO:3 to denote degenerate nucleotide positions. "Resolutions" are the nucleotides denoted by a code letter. "Complement" indicates the code for the complementary nucleotide(s). For example, the code Y denotes either C or T, and its complement R denotes A or G, A
30 being complementary to T, and G being complementary to C.

Table 1

Nucleotide	Resolution	Complement	Resolution
A	A	T	T
C	C	C	G
G	C	C	C
T	T	A	A
R	A G	B	C T
Y	C T	B	A G
M	A G	B	G T
K	G T	M	A C
S	C G	S	C G
W	A T	W	A T
H	A C T	D	A G T
B	C G T	V	A C G
V	A C G	B	C G T
D	A C T	H	A C T
N	A C G T	N	A C G T

The degenerate codons used in SEQ ID NO:3, encompassing all possible
5 codons for a given amino acid, are set forth in Table 2.

Table 2

Amino Acid	One Letter Code	Codons	Degenerate Codon
Cys	C	TGC TGT	TTY
Ser	S	AGC AGT TCA TCC TCG TCT	WSN
Thr	T	ACA ACC ACG ACT	ACN
Pro	P	CCA CCC CCG CCT	CCN
Ala	A	GCA GCC GCG GCT	GCN
Gly	G	GGA GGC GGG GGT	GGN
Asn	N	AAC AAT	AAY
Asp	D	GAC GAT	GAY
Glu	L	GAA GAG	GAR
Gln	Q	CAA CAG	CAR
His	H	CAC CAT	CAY
Arg	R	AGA AGG CGA CGC CGG CGT	MGN
Lys	K	AAA AAG	AAR
Met	M	ATG	ATG
Ile	I	ATA ATC ATT	ATH
Leu	L	CTA CTC CTG CTT TTA TTG	YTN
Val	V	GTA GTC GTG GTT	GTN
Phe	F	TTC TTT	TTY
Tyr	Y	TAC TAT	TAY
Trp	W	TGG	TGG
Ter	.	TAA TAG TGA	TRR
Asn Asp	B		RAY
Glu Gln	Z		SAR
Any	X		NNN

One of ordinary skill in the art will appreciate that some ambiguity is introduced in determining a degenerate codon, representative of all possible codons encoding an amino acid. For example, the degenerate codon for serine (WSN) can, in some circumstances, encode arginine (AGR), and the degenerate codon for arginine (MGN) can, in some circumstances, encode serine (AGY). A similar relationship exists between codons encoding phenylalanine and leucine. Thus, some polynucleotides encompassed by the degenerate sequence may encode variant amino acid sequences, but one of ordinary skill in the art can easily identify such variant sequences by reference to the amino acid sequence of SEQ ID NO:2. Variant sequences can be readily tested for functionality as described herein.

Different species can exhibit "preferential codon usage." In general, see, Grantham *et al.*, *Nuc. Acids Res.* 8:1893 (1980), Haas *et al.* *Curr. Biol.* 6:315 (1996), Wain-Hobson *et al.*, *Gene* 13:355 (1981), Grosjean and Fiers, *Gene* 18:199 (1982), Holm, *Nuc. Acids Res.* 14:3075 (1986), Ikemura, *J. Mol. Biol.* 158:573 (1982), Sharp and Matassi, *Curr. Opin. Genet. Dev.* 4:851 (1994), Kane, *Curr. Opin. Biotechnol.* 6:494 (1995), and Makrides, *Microbiol. Rev.* 60:512 (1996). As used herein, the term "preferential codon usage" or "preferential codons" is a term of art referring to protein translation codons that are most frequently used in cells of a certain species, thus favoring one or a few representatives of the possible codons encoding each amino acid (see Table 2). For example, the amino acid threonine (thr) may be encoded by ACA, ACC, ACG, or ACT, but in mammalian cells ACC is the most commonly used codon; in other species, for example, insect cells, yeast, viruses or bacteria, different thr codons may be preferential. Preferential codons for a particular species can be introduced into the polynucleotides of the present invention by a variety of methods known in the art. Introduction of preferential codon sequences into recombinant DNA can, for example, enhance production of the protein by making protein translation more efficient within a particular cell type or species. Therefore, the degenerate codon sequence disclosed in SEQ ID NO:3 serves as a template for optimizing expression of polynucleotides in various cell types and species commonly used in the art and disclosed herein. Sequences containing preferential codons can be tested and optimized for expression in various species, and tested for functionality as disclosed herein.

The present invention further provides variant polypeptides and nucleic acid molecules that represent counterparts from other species (orthologs). These species include, but are not limited to mammalian, avian, amphibian, reptile, fish, insect and other vertebrate and invertebrate species. Of particular interest are Zmim19 polypeptides from other mammalian species, including porcine, murine, ovine, bovine, canine, feline, equine, and other primate polypeptides. As an illustration, SEQ ID

NO:5, SEQ ID NO:6, and SEQ ID NO:7 provide the nucleotide, amino acid, and degenerate nucleotide sequences, respectively, of murine Zmim19. Sequence analysis indicates that the murine Zmim19 polypeptide includes active site residues at His²⁰⁷, Cys²⁵⁹, Asp²⁹², and Asp²⁹³.

5 Orthologs of human Zmim19 can be cloned using information and compositions provided by the present invention in combination with conventional cloning techniques. For example, a cDNA can be cloned using mRNA obtained from a tissue or cell type that expresses Zmim19 as disclosed herein. Suitable sources of mRNA can be identified by probing northern blots with probes designed from the
10 sequences disclosed herein. A library is then prepared from mRNA of a positive tissue or cell line.

A Zmim19-encoding cDNA can then be isolated by a variety of methods, such as by probing with a complete or partial human cDNA or with one or more sets of degenerate probes based on the disclosed sequences. A cDNA can also be
15 cloned using the polymerase chain reaction with primers designed from the representative human Zmim19 sequences disclosed herein. Within an additional method, the cDNA library can be used to transform or transfect host cells, and expression of the cDNA of interest can be detected with an antibody to Zmim19 polypeptide. Similar techniques can also be applied to the isolation of genomic clones.

20 Those skilled in the art will recognize that the sequence disclosed in SEQ ID NO:1 represents a single allele of human Zmim19, and that allelic variation and alternative splicing are expected to occur. Allelic variants of this sequence can be cloned by probing cDNA or genomic libraries from different individuals according to standard procedures. Allelic variants of the nucleotide sequence shown in SEQ ID
25 NO:1, including those containing silent mutations and those in which mutations result in amino acid sequence changes, are within the scope of the present invention, as are proteins which are allelic variants of SEQ ID NO:2. cDNA molecules generated from alternatively spliced mRNAs, which retain the properties of the Zmim19 polypeptide are included within the scope of the present invention, as are polypeptides encoded by
30 such cDNAs and mRNAs. Allelic variants and splice variants of these sequences can be cloned by probing cDNA or genomic libraries from different individuals or tissues according to standard procedures known in the art.

Within certain embodiments of the invention, the isolated nucleic acid molecules can hybridize under stringent conditions to nucleic acid molecules
35 comprising nucleotide sequences disclosed herein. For example, such nucleic acid molecules can hybridize under stringent conditions to nucleic acid molecules comprising the nucleotide sequence of nucleotides 172 to 1236 of SEQ ID NO:1, to

nucleic acid molecules consisting of the nucleotide sequence of SEQ ID NO:1, or to nucleic acid molecules consisting of a nucleotide sequence complementary to nucleotides 172 to 1236 of SEQ ID NO:1, or to SEQ ID NO:1. In general, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe.

A pair of nucleic acid molecules, such as DNA-DNA, RNA-RNA and DNA-RNA, can hybridize if the nucleotide sequences have some degree of complementarity. Hybrids can tolerate mismatched base pairs in the double helix, but the stability of the hybrid is influenced by the degree of mismatch. The T_m of the mismatched hybrid decreases by 1°C for every 1-1.5% base pair mismatch. Varying the stringency of the hybridization conditions allows control over the degree of mismatch that will be present in the hybrid. The degree of stringency increases as the hybridization temperature increases and the ionic strength of the hybridization buffer decreases. Stringent hybridization conditions encompass temperatures of about 5-25°C below the T_m of the hybrid and a hybridization buffer having up to 1 M Na^+ . Higher degrees of stringency at lower temperatures can be achieved with the addition of formamide which reduces the T_m of the hybrid about 1°C for each 1% formamide in the buffer solution. Generally, such stringent conditions include temperatures of 20-70°C and a hybridization buffer containing up to 6x SSC and 0-50% formamide. A higher degree of stringency can be achieved at temperatures of from 40-70°C with a hybridization buffer having up to 4x SSC and from 0-50% formamide. Highly stringent conditions typically encompass temperatures of 42-70°C with a hybridization buffer having up to 1x SSC and 0-50% formamide. Different degrees of stringency can be used during hybridization and washing to achieve maximum specific binding to the target sequence. Typically, the washes following hybridization are performed at increasing degrees of stringency to remove non-hybridized polynucleotide probes from hybridized complexes.

The above conditions are meant to serve as a guide and it is well within the abilities of one skilled in the art to adapt these conditions for use with a particular polypeptide hybrid. The T_m for a specific target sequence is the temperature (under defined conditions) at which 50% of the target sequence will hybridize to a perfectly matched probe sequence. Those conditions that influence the T_m include, the size and base pair content of the polynucleotide probe, the ionic strength of the hybridization solution, and the presence of destabilizing agents in the hybridization solution. Numerous equations for calculating T_m are known in the art, and are specific for DNA,

RNA and DNA-RNA hybrids and polynucleotide probe sequences of varying length (see, for example, Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Second Edition (Cold Spring Harbor Press 1989); Ausubel *et al.*, (eds.), *Current Protocols in Molecular Biology* (John Wiley and Sons, Inc. 1987); Berger and Kimmel (eds.), *Guide to Molecular Cloning Techniques*, (Academic Press, Inc. 1987); and Wetmur, *Crit. Rev. Biochem. Mol. Biol.* 26:227 (1990)). Sequence analysis software such as OLIGO 6.0 (LSR; Long Lake, MN) and *Primer Premier 4.0* (Premier Biosoft International; Palo Alto, CA), as well as sites on the Internet, are available tools for analyzing a given sequence and calculating T_m based on user defined criteria. Such programs can also analyze a given sequence under defined conditions and identify suitable probe sequences. Typically, hybridization of longer polynucleotide sequences, >50 base pairs, is performed at temperatures of about 20-25°C below the calculated T_m . For smaller probes, <50 base pairs, hybridization is typically carried out at the T_m or 5-10°C below. This allows for the maximum rate of hybridization for DNA-DNA and DNA-RNA hybrids.

The length of the polynucleotide sequence influences the rate and stability of hybrid formation. Smaller probe sequences, <50 base pairs, reach equilibrium with complementary sequences rapidly, but may form less stable hybrids. Incubation times of anywhere from minutes to hours can be used to achieve hybrid formation. Longer probe sequences come to equilibrium more slowly, but form more stable complexes even at lower temperatures. Incubations are allowed to proceed overnight or longer. Generally, incubations are carried out for a period equal to three times the calculated Cot time. Cot time, the time it takes for the polynucleotide sequences to reassociate, can be calculated for a particular sequence by methods known in the art.

The base pair composition of polynucleotide sequence will effect the thermal stability of the hybrid complex, thereby influencing the choice of hybridization temperature and the ionic strength of the hybridization buffer. A-T pairs are less stable than G-C pairs in aqueous solutions containing sodium chloride. Therefore, the higher the G-C content, the more stable the hybrid. Even distribution of G and C residues within the sequence also contribute positively to hybrid stability. In addition, the base pair composition can be manipulated to alter the T_m of a given sequence. For example, 5-methyldeoxycytidine can be substituted for deoxycytidine and 5-bromodeoxuridine can be substituted for thymidine to increase the T_m , whereas 7-deaz-2'-deoxyguanosine can be substituted for guanosine to reduce dependence on T_m .

The ionic concentration of the hybridization buffer also affects the stability of the hybrid. Hybridization buffers generally contain blocking agents such as

Denhardt's solution (Sigma Chemical Co., St. Louis, Mo.), denatured salmon sperm DNA, tRNA, milk powders (BLOTTO), heparin or SDS, and a Na^+ source, such as SSC (1x SSC: 0.15 M sodium chloride, 15 mM sodium citrate) or SSPE (1x SSPE: 1.8 M NaCl, 10 mM NaH_2PO_4 , 1 mM EDTA, pH 7.7). By decreasing the ionic concentration of the buffer, the stability of the hybrid is increased. Typically, hybridization buffers contain from between 10 mM - 1 M Na^+ . The addition of destabilizing or denaturing agents such as formamide, tetralkylammonium salts, guanidinium cations or thiocyanate cations to the hybridization solution will alter the T_m of a hybrid. Typically, formamide is used at a concentration of up to 50% to allow incubations to be carried out at more convenient and lower temperatures. Formamide also acts to reduce non-specific background when using RNA probes.

As an illustration, a nucleic acid molecule encoding a variant Zn_{mim}19 polypeptide can be hybridized with a nucleic acid molecule having the nucleotide sequence of nucleotides 172 to 1236 of SEQ ID NO:1 (or its complement) at 42°C overnight in a solution comprising 50% formamide, 5xSSC (1xSSC: 0.15 M sodium chloride and 15 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution (100x Denhardt's solution: 2% (w/v) Ficoll 400, 2% (w/v) polyvinylpyrrolidone, and 2% (w/v) bovine serum albumin, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA. One of skill in the art can devise variations of these hybridization conditions. For example, the hybridization mixture can be incubated at a higher temperature, such as about 65°C, in a solution that does not contain formamide. Moreover, premixed hybridization solutions are available (e.g., EXPRESSHYB Hybridization Solution from CLONTECH Laboratories, Inc.), and hybridization can be performed according to the manufacturer's instructions.

Following hybridization, the nucleic acid molecules can be washed to remove non-hybridized nucleic acid molecules under stringent conditions, or under highly stringent conditions. Typical stringent washing conditions include washing in a solution of 0.5x - 2x SSC with 0.1% sodium dodecyl sulfate (SDS) at 55 - 65°C. As an illustration, particular nucleic acid molecules encoding a variant Zn_{mim}19 polypeptide remain hybridized following stringent washing conditions with a nucleic acid molecule consisting of the nucleotide sequence of nucleotides 172 to 1236 of SEQ ID NO:1 (or its complement), in which the wash stringency is equivalent to 0.5x - 2x SSC with 0.1% SDS at 55 - 65°C, including 0.5x SSC with 0.1% SDS at 55°C, or 2xSSC with 0.1% SDS at 65°C. One of skill in the art can readily devise equivalent conditions, for example, by substituting the SSPE for SSC in the wash solution.

Typical highly stringent washing conditions include washing in a solution of 0.1x - 0.2x SSC with 0.1% sodium dodecyl sulfate (SDS) at 50 - 65°C. For

example, certain nucleic acid molecules encoding a variant Zmim19 polypeptide remain hybridized following highly stringent washing conditions with a nucleic acid molecule consisting of the nucleotide sequence of nucleotides 172 to 1236 of SEQ ID NO:1 (or its complement), in which the wash stringency is equivalent to 0.1x - 0.2x SSC with 0.1% SDS at 50 - 65°C, including 0.1x SSC with 0.1% SDS at 50°C, or 0.2xSSC with 0.1% SDS at 65°C.

The present invention also provides isolated Zmim19 polypeptides that have a substantially similar sequence identity to the polypeptide of SEQ ID NO:2, or orthologs. The term "substantially similar sequence identity" is used herein to denote polypeptides having 70%, 80%, 90%, 95%, 96%, 97%, 98%, or 99% sequence identity to the sequence shown in SEQ ID NO:2.

The present invention also contemplates Zmim19 variant nucleic acid molecules that can be identified using two criteria: a determination of the similarity between the encoded polypeptide with the amino acid sequence of SEQ ID NO:2, and a hybridization assay, as described above. Such Zmim19 variants include nucleic acid molecules (1) that remain hybridized following stringent washing conditions with a nucleic acid molecule consisting of the nucleotide sequence of nucleotides 172 to 1236 of SEQ ID NO:1 (or its complement), in which the wash stringency is equivalent to 0.5x - 2x SSC with 0.1% SDS at 55 - 65°C, and (2) that encode a polypeptide having 70%, 80%, 90%, 95% 96%, 97%, 98% or 99% sequence identity to the amino acid sequence of SEQ ID NO:2.

Alternatively, Zmim19 variants can be characterized as nucleic acid molecules (1) that remain hybridized following highly stringent washing conditions with a nucleic acid molecule consisting of the nucleotide sequence of nucleotides 172 to 1236 of SEQ ID NO:1 (or its complement), in which the wash stringency is equivalent to 0.1x - 0.2x SSC with 0.1% SDS at 50 - 65°C, and (2) that encode a polypeptide having 70%, 80%, 90%, 95%, 96%, 97%, 98% or 99% sequence identity to the amino acid sequence of SEQ ID NO:2.

The present invention also includes Zmim19 variants that possess adenosine deaminase enzyme activity. Moreover, particular Zmim19 variants are characterized using hybridization analysis with a reference nucleic acid molecule that is a fragment of a nucleic acid molecule consisting of the nucleotide sequence of nucleotides 172 to 1236 of SEQ ID NO:1, or its complement.

Percent sequence identity is determined by conventional methods. See, for example, Altschul *et al.*, *Bull. Math. Bio.* 48:603 (1986), and Henikoff and Henikoff, *Proc. Nat'l Acad. Sci. USA* 89:10915 (1992). Briefly, two amino acid sequences are aligned to optimize the alignment scores using a gap opening penalty of

10, a gap extension penalty of 1, and the “BLOSUM62” scoring matrix of Henikoff and Henikoff (*ibid.*) as shown in Table 3 (amino acids are indicated by the standard one-letter codes). The percent identity is then calculated as: ([Total number of identical matches]/ [length of the longer sequence plus the number of gaps introduced into the
5 longer sequence in order to align the two sequences])(100).

Table 3

	A	R	N	D	C	Q	E	G	H	I	L	K	M	P	S	T	W	Y		
5	A	4																		
	R	-1	5																	
	N	-2	0	6																
	D	-2	-2	1	6															
	C	0	-3	-3	-3	9														
	Q	-1	1	0	-3	5														
	E	-1	0	0	-4	2	5													
	G	0	-2	0	-1	-3	-2	-2	6											
	H	-2	0	1	-1	-3	0	0	-2	8										
	I	-1	-3	-3	-1	-3	-3	-4	-3	4										
	L	-1	-2	-3	-4	-1	-2	-3	-4	-3	2	4								
	X	-1	2	0	-1	-3	1	1	-2	-1	-3	-2	5							
	M	-1	-1	-2	-3	-1	0	-2	-3	-2	1	2	-1	5						
	P	-2	-3	-3	-2	-3	-3	-1	0	-3	0	6								
	F	-1	-2	-2	-1	-3	-1	-1	-2	-2	-3	-1	-2	-4	7					
	S	-1	-1	0	-1	0	0	0	-1	-2	-2	0	-1	-2	-1	4				
	T	0	-1	0	-1	-1	-1	-1	-2	-2	-1	-1	-1	-2	-1	1	5			
	W	-3	-3	-4	-4	-2	-2	-3	-2	-2	-3	-3	-1	-4	-3	-2	11			
	Y	-2	-2	-2	-2	-1	-2	-3	2	-1	-2	-1	3	-3	-2	2	7			
	V	0	-3	-3	-1	-2	-2	-3	-3	3	1	-2	1	-1	-2	2	0	-3	-1	4

Those skilled in the art appreciate that there are many established algorithms available to align two amino acid sequences. The "FASTA" similarity search algorithm of Pearson and Lipman is a suitable protein alignment method for examining the level of identity shared by an amino acid sequence disclosed herein and the amino acid sequence of a putative Zmim19 variant. The FASTA algorithm is described by Pearson and Lipman, *Proc. Nat'l Acad. Sci. USA* 85:2444 (1988), and by Pearson, *Meth. Enzymol.* 183:63 (1990). Briefly, FASTA first characterizes sequence similarity by identifying regions shared by the query sequence (e.g., SEQ ID NO:2) and a test sequence that have either the highest density of identities (if the ktup variable is 1) or pairs of identities (if ktup=2), without considering conservative amino acid substitutions, insertions, or deletions. The ten regions with the highest density of identities are then rescored by comparing the similarity of all paired amino acids using an amino acid substitution matrix, and the ends of the regions are "trimmed" to include only those residues that contribute to the highest score. If there are several regions with scores greater than the "cutoff" value (calculated by a predetermined formula based upon the length of the sequence and the ktup value), then the trimmed initial regions are examined to determine whether the regions can be joined to form an approximate alignment with gaps. Finally, the highest scoring regions of the two amino acid sequences are aligned using a modification of the Needleman-Wunsch-Sellers algorithm (Needleman and Wunsch, *J. Mol. Biol.* 48:444 (1970); Sellers, *SIAM J. Appl. Math.* 26:787 (1974)), which allows for amino acid insertions and deletions. Illustrative parameters for FASTA analysis are: ktup=1, gap opening penalty=10, gap extension penalty=1, and substitution matrix=BLOSUM62. These parameters can be introduced into a FASTA program by modifying the scoring matrix file ("SMATRIX"), as explained in Appendix 2 of Pearson, *Meth. Enzymol.* 183:63 (1990).

FASTA can also be used to determine the sequence identity of nucleic acid molecules using a ratio as disclosed above. For nucleotide sequence comparisons, the ktup value can range between one to six, preferably from three to six, most preferably three, with other parameters set as described above.

The present invention includes nucleic acid molecules that encode a polypeptide having a conservative amino acid change, compared with the amino acid sequence of SEQ ID NO:2. That is, variants can be obtained that contain one or more amino acid substitutions of SEQ ID NO:2, in which an alkyl amino acid is substituted for an alkyl amino acid in a Zmim19 amino acid sequence, an aromatic amino acid is substituted for an aromatic amino acid in a Zmim19 amino acid sequence, a sulfur-containing amino acid is substituted for a sulfur-containing amino acid in a Zmim19 amino acid sequence, a hydroxy-containing amino acid is substituted for a hydroxy-

containing amino acid in a Zmim19 amino acid sequence, an acidic amino acid is substituted for an acidic amino acid in a Zmim19 amino acid sequence, a basic amino acid is substituted for a basic amino acid in a Zmim19 amino acid sequence, or a dibasic monocarboxylic amino acid is substituted for a dibasic monocarboxylic amino acid in a Zmim19 amino acid sequence.

Among the common amino acids, for example, a "conservative amino acid substitution" is illustrated by a substitution among amino acids within each of the following groups: (1) glycine, alanine, valine, leucine, and isoleucine, (2) phenylalanine, tyrosine, and tryptophan, (3) serine and threonine, (4) aspartate and glutamate, (5) glutamine and asparagine, and (6) lysine, arginine and histidine.

The BLOSUM62 table is an amino acid substitution matrix derived from about 2,000 local multiple alignments of protein sequence segments, representing highly conserved regions of more than 500 groups of related proteins (Henikoff and Henikoff, *Proc. Nat'l Acad. Sci. USA* 89:10915 (1992)). Accordingly, the BLOSUM62 substitution frequencies can be used to define conservative amino acid substitutions that may be introduced into the amino acid sequences of the present invention. Although it is possible to design amino acid substitutions based solely upon chemical properties (as discussed above), the language "conservative amino acid substitution" preferably refers to a substitution represented by a BLOSUM62 value of greater than -1. For example, an amino acid substitution is conservative if the substitution is characterized by a BLOSUM62 value of 0, 1, 2, or 3. According to this system, preferred conservative amino acid substitutions are characterized by a BLOSUM62 value of at least 1 (e.g., 1, 2 or 3), while more preferred conservative amino acid substitutions are characterized by a BLOSUM62 value of at least 2 (e.g., 2 or 3).

Particular variants of Zmim19 are characterized by having greater than 96%, at least 97%, at least 98%, or at least 99% sequence identity to the corresponding amino acid sequence (e.g., SEQ ID NO:2), wherein the variation in amino acid sequence is due to one or more conservative amino acid substitutions.

Conservative amino acid changes in a Zmim19 gene can be introduced by substituting nucleotides for the nucleotides recited in SEQ ID NO:1. Such "conservative amino acid" variants can be obtained, for example, by oligonucleotide-directed mutagenesis, linker-scanning mutagenesis, mutagenesis using the polymerase chain reaction, and the like (see Ausubel (1995) at pages 8-10 to 8-22; and McPherson (ed.), *Directed Mutagenesis: A Practical Approach* (IRL Press 1991)).

The proteins of the present invention can also comprise non-naturally occurring amino acid residues. Non-naturally occurring amino acids include, without limitation, *trans*-3-methylproline, 2,4-methanoproline, *cis*-4-hydroxyproline, *trans*-4-

hydroxyproline, *N*-methylglycine, *allo*-threonine, methylthreonine, hydroxyethylcysteine, hydroxyethylhomocysteine, nitroglutamine, homoglutamine, pipecolic acid, thiazolidine carboxylic acid, dehydropipecolic acid, 3- and 4-methylproline, 3,3-dimethylproline, *tert*-leucine, norvaline, 2-azaphenylalanine, 3-azaphenylalanine, 4-azaphenylalanine, and 4-fluorophenylalanine. Several methods are known in the art for incorporating non-naturally occurring amino acid residues into proteins. For example, an *in vitro* system can be employed wherein nonsense mutations are suppressed using chemically aminoacylated suppressor tRNAs. Methods for synthesizing amino acids and aminoacylating tRNA are known in the art. Transcription and translation of plasmids containing nonsense mutations is typically carried out in a cell-free system comprising an *E. coli* S30 extract and commercially available enzymes and other reagents. Proteins are purified by chromatography. See, for example, Robertson *et al.*, *J. Am. Chem. Soc.* 113:2722 (1991), Ellman *et al.*, *Methods Enzymol.* 202:301 (1991), Chung *et al.*, *Science* 259:806 (1993), and Chung *et al.*, *Proc. Nat'l Acad. Sci. USA* 90:10145 (1993).

In a second method, translation is carried out in *Xenopus* oocytes by microinjection of mutated mRNA and chemically aminoacylated suppressor tRNAs (Turcatti *et al.*, *J. Biol. Chem.* 271:19991 (1996)). Within a third method, *E. coli* cells are cultured in the absence of a natural amino acid that is to be replaced (e.g., phenylalanine) and in the presence of the desired non-naturally occurring amino acid(s) (e.g., 2-azaphenylalanine, 3-azaphenylalanine, 4-azaphenylalanine, or 4-fluorophenylalanine). The non-naturally occurring amino acid is incorporated into the protein in place of its natural counterpart. See, Koide *et al.*, *Biochem.* 33:7470 (1994). Naturally occurring amino acid residues can be converted to non-naturally occurring species by *in vitro* chemical modification. Chemical modification can be combined with site-directed mutagenesis to further expand the range of substitutions (Wynn and Richards, *Protein Sci.* 2:395 (1993)).

A limited number of non-conservative amino acids, amino acids that are not encoded by the genetic code, non-naturally occurring amino acids, and unnatural amino acids may be substituted for Zmim19 amino acid residues.

Essential amino acids in the polypeptides of the present invention can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, *Science* 244:1081 (1989), Bass *et al.*, *Proc. Nat'l Acad. Sci. USA* 88:4498 (1991), Coombs and Corey, "Site-Directed Mutagenesis and Protein Engineering," in *Proteins: Analysis and Design*, Angeletti (ed.), pages 259-311 (Academic Press, Inc. 1998)). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for biological activity as disclosed below to identify amino acid

residues that are critical to the activity of the molecule. See also, Hilton *et al.*, *J. Biol. Chem.* 271:4699 (1996). The identities of essential amino acids can also be inferred from analysis of homologies with other adenosine deaminases.

The location of Zmim19 activity domains can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos *et al.*, *Science* 255:306 (1992), Smith *et al.*, *J. Mol. Biol.* 224:899 (1992), and Wlodaver *et al.*, *FEBS Lett.* 309:59 (1992).

Multiple amino acid substitutions can be made and tested using known methods of mutagenesis and screening, such as those disclosed by Reidhaar-Olson and Sauer (*Science* 241:53 (1988)) or Bowie and Sauer (*Proc. Nat'l Acad. Sci. USA* 86:2152 (1989)). Briefly, these authors disclose methods for simultaneously randomizing two or more positions in a polypeptide, selecting for functional polypeptide, and then sequencing the mutagenized polypeptides to determine the spectrum of allowable substitutions at each position. Other methods that can be used include phage display (e.g., Lowman *et al.*, *Biochem.* 30:10832 (1991), Ladner *et al.*, U.S. Patent No. 5,223,409, Huse, international publication No. WO 92/06204, and region-directed mutagenesis (Derbyshire *et al.*, *Gene* 46:145 (1986), and Ner *et al.*, *DNA* 7:127, (1988)).

Variants of the disclosed Zmim19 nucleotide and polypeptide sequences can also be generated through DNA shuffling as disclosed by Stemmer, *Nature* 370:389 (1994), Stemmer, *Proc. Nat'l Acad. Sci. USA* 91:10747 (1994), and international publication No. WO 97/20078. Briefly, variant DNAs are generated by *in vitro* homologous recombination by random fragmentation of a parent DNA followed by reassembly using PCR, resulting in randomly introduced point mutations. This technique can be modified by using a family of parent DNAs, such as allelic variants or DNAs from different species, to introduce additional variability into the process. Selection or screening for the desired activity, followed by additional iterations of mutagenesis and assay provides for rapid "evolution" of sequences by selecting for desirable mutations while simultaneously selecting against detrimental changes.

Mutagenesis methods as disclosed herein can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides in host cells. Mutagenized DNA molecules that encode biologically active polypeptides, or polypeptides that bind with anti-Zmim19 antibodies, can be recovered from the host cells and rapidly sequenced using modern equipment. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

The present invention also includes "functional fragments" of Zmim19 polypeptides and nucleic acid molecules encoding such functional fragments. Routine deletion analyses of nucleic acid molecules can be performed to obtain functional fragments of a nucleic acid molecule that encodes a Zmim19 polypeptide. As an illustration, DNA molecules having the nucleotide sequence of SEQ ID NO:1 can be digested with *Bal31* nuclease to obtain a series of nested deletions. The fragments are then inserted into expression vectors in proper reading frame, and the expressed polypeptides are isolated and tested for adenosine deaminase activity. One alternative to exonuclease digestion is to use oligonucleotide-directed mutagenesis to introduce deletions or stop codons to specify production of a desired fragment. Alternatively, particular fragments of a *Zmim19* gene can be synthesized using the polymerase chain reaction.

As an illustration of this general approach, studies on the truncation at either or both termini of interferons have been summarized by Horisberger and Di Marco, *Pharmac. Ther.* 66:507 (1995). Moreover, standard techniques for functional analysis of proteins are described by, for example, Treuter *et al.*, *Molec. Gen. Genet.* 240:113 (1993), Content *et al.*, "Expression and preliminary deletion analysis of the 42 kDa 2-5A synthetase induced by human interferon," in *Biological Interferon Systems, Proceedings of ISIR-TNO Meeting on Interferon Systems*, Cantell (ed.), pages 65-72 (Nijhoff 1987), Herschman, "The EGF Receptor," in *Control of Animal Cell Proliferation, Vol. 1*, Boynton *et al.*, (eds.) pages 169-199 (Academic Press 1985), Coumilleau *et al.*, *J. Biol. Chem.* 270:29270 (1995); Fukunaga *et al.*, *J. Biol. Chem.* 270:25291 (1995); Yamaguchi *et al.*, *Biochem. Pharmacol.* 50:1295 (1995), and Meisel *et al.*, *Plant Molec. Biol.* 30:1 (1996).

The present invention also contemplates functional fragments of a *Zmim19* gene that has amino acid changes, compared with the amino acid sequence of SEQ ID NO:2. A variant *Zmim19* gene can be identified on the basis of structure by determining the level of identity with nucleotide and amino acid sequences of SEQ ID NOs:1 and 2, as discussed above. An alternative approach to identifying a variant gene on the basis of structure is to determine whether a nucleic acid molecule encoding a potential variant *Zmim19* gene can hybridize to a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1, as discussed above.

The present invention also provides polypeptide fragments or peptides comprising an epitope-bearing portion of a *Zmim19* polypeptide described herein. Such fragments or peptides may comprise an "immunogenic epitope," which is a part of a protein that elicits an antibody response when the entire protein is used as an

immunogen. Immunogenic epitope-bearing peptides can be identified using standard methods (see, for example, Geysen *et al.*, *Proc. Nat'l Acad. Sci. USA* 81:3998 (1983)).

In contrast, polypeptide fragments or peptides may comprise an "antigenic epitope," which is a region of a protein molecule to which an antibody can specifically bind. Certain epitopes consist of a linear or contiguous stretch of amino acids, and the antigenicity of such an epitope is not disrupted by denaturing agents. It is known in the art that relatively short synthetic peptides that can mimic epitopes of a protein can be used to stimulate the production of antibodies against the protein (see, for example, Sutcliffe *et al.*, *Science* 219:660 (1983)). Antibodies that recognize short linear epitopes are particularly useful in analytic and diagnostic applications that use denatured protein, such as Western analysis, or in the analysis of fixed cells or tissue samples. Antibodies to linear epitopes are also useful for detecting fragments of Zmin19, such as might occur in body fluids or culture media. Accordingly, antigenic epitope-bearing peptides and polypeptides of the present invention are useful to raise antibodies that bind with the polypeptides described herein.

Antigenic epitope-bearing peptides and polypeptides can contain at least four to ten amino acids, at least ten to fifteen amino acids, or about 15 to about 30 amino acids of SEQ ID NO:2. Such epitope-bearing peptides and polypeptides can be produced by fragmenting a Zmin19 polypeptide, or by chemical peptide synthesis, as described herein. Moreover, epitopes can be selected by phage display of random peptide libraries (see, for example, Lane and Stephen, *Curr. Opin. Immunol.* 5:268 (1993), and Cortese *et al.*, *Curr. Opin. Biotechnol.* 7:616 (1996)). Standard methods for identifying epitopes and producing antibodies from small peptides that comprise an epitope are described, for example, by Mole, "Epitope Mapping," in *Methods in Molecular Biology*, Vol. 10, Manson (ed.), pages 105-116 (The Humana Press, Inc. 1992), Price, "Production and Characterization of Synthetic Peptide-Derived Antibodies," in *Monoclonal Antibodies: Production, Engineering, and Clinical Application*, Ritter and Ladyman (eds.), pages 60-84 (Cambridge University Press 1995), and Coligan *et al.* (eds.), *Current Protocols in Immunology*, pages 9.3.1 - 9.3.5 and pages 9.4.1 - 9.4.11 (John Wiley & Sons 1997).

For any Zmin19 polypeptide, including variants and fusion proteins, one of ordinary skill in the art can readily generate a fully degenerate polynucleotide sequence encoding that variant using the information set forth in Tables 1 and 2 above. Moreover, those of skill in the art can use standard software to devise Zmin19 variants based upon the nucleotide and amino acid sequences described herein. Accordingly, the present invention includes a computer-readable medium encoded with a data structure that provides at least one of SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3. Suitable forms of computer-readable media include magnetic media and optically-readable media.

Examples of magnetic media include a hard or fixed drive, a random access memory (RAM) chip, a floppy disk, digital linear tape (DLT), a disk cache, and a ZIP disk. Optically readable media are exemplified by compact discs (*e.g.*, CD-read only memory (ROM), CD-rewritable (RW), and CD-recordable), and digital versatile/video discs (DVD) (*e.g.*, DVD-ROM, DVD-RAM, and DVD+RW).

5. **Production of Zmim19 Fusion Proteins**

Fusion proteins of Zmim19 can be used to express Zmim19 in a recombinant host, and to isolate expressed Zmim19. One type of fusion protein comprises a peptide that guides a Zmim19 polypeptide from a recombinant host cell. To direct a Zmim19 polypeptide into the secretory pathway of a eukaryotic host cell, a secretory signal sequence (also known as a signal peptide, a leader sequence, prepro sequence or pre sequence) is provided in the Zmim19 expression vector. A suitable signal sequence may be derived from a secreted protein or synthesized *de novo*. The secretory signal sequence is operably linked to a Zmim19-encoding sequence such that the two sequences are joined in the correct reading frame and positioned to direct the newly synthesized polypeptide into the secretory pathway of the host cell. Secretory signal sequences are commonly positioned 5' to the nucleotide sequence encoding the polypeptide of interest, although certain secretory signal sequences may be positioned elsewhere in the nucleotide sequence of interest (see, *e.g.*, Welch *et al.*, U.S. Patent No. 5,037,743; Holland *et al.*, U.S. Patent No. 5,143,830).

While the secretory signal sequence of a protein produced by mammalian cells (*e.g.*, tissue-type plasminogen activator signal sequence, as described, for example, in U.S. Patent No. 5,641,655) is useful for expression of Zmim19 in recombinant mammalian hosts, a yeast signal sequence is preferred for expression in yeast cells. Examples of suitable yeast signal sequences are those derived from yeast mating pheromone α -factor (encoded by the *MF α* gene), invertase (encoded by the *SUC2* gene), or acid phosphatase (encoded by the *PHO5* gene). See, for example, Romanos *et al.*, "Expression of Cloned Genes in Yeast," in *DNA Cloning 2: A Practical Approach*, 2nd Edition, Glover and Hames (eds.), pages 123-167 (Oxford University Press 1995).

In bacterial cells, it is often desirable to express a heterologous protein as a fusion protein to decrease toxicity, increase stability, and to enhance recovery of the expressed protein. For example, Zmim19 can be expressed as a fusion protein comprising a glutathione S-transferase polypeptide. Glutathione S-transferase fusion proteins are typically soluble, and easily purifiable from *E. coli* lysates on immobilized glutathione columns. In similar approaches, a Zmim19 fusion protein comprising a

maltose binding protein polypeptide can be isolated with an amylose resin column, while a fusion protein comprising the C-terminal end of a truncated Protein A gene can be purified using IgG-Sepharose. Established techniques for expressing a heterologous polypeptide as a fusion protein in a bacterial cell are described, for example, by Williams *et al.*, "Expression of Foreign Proteins in *E. coli* Using Plasmid Vectors and Purification of Specific Polyclonal Antibodies," in *DNA Cloning 2: A Practical Approach*, 2nd Edition, Glover and Hames (Eds.), pages 15-58 (Oxford University Press 1995). In addition, commercially available expression systems are available. For example, the PINPOINT Xa protein purification system (Promega Corporation; Madison, WI) provides a method for isolating a fusion protein comprising a polypeptide that becomes biotinylated during expression with a resin that comprises avidin.

Peptide tags that are useful for isolating heterologous polypeptides expressed by either prokaryotic or eukaryotic cells include polyHistidine tags (which have an affinity for nickel-chelating resin), *c-myc* tags, calmodulin binding protein (isolated with calmodulin affinity chromatography), substance P, the RYRS tag (which binds with anti-RYRS antibodies), the Glu-Glu tag, and the FLAG tag (which binds with anti-FLAG antibodies). See, for example, Luo *et al.*, *Arch. Biochem. Biophys.* 329:215 (1996), Morganti *et al.*, *Biotechnol. Appl. Biochem.* 23:67 (1996), and Zheng *et al.*, *Gene* 186:55 (1997). Nucleic acid molecules encoding such peptide tags are available, for example, from Sigma-Aldrich Corporation (St. Louis, MO).

Another form of fusion protein comprises a Zmim19 polypeptide and an immunoglobulin heavy chain constant region, typically an F_c fragment, which contains two constant region domains and a hinge region but lacks the variable region. As an illustration, Chang *et al.*, U.S. Patent No. 5,723,125, describe a fusion protein comprising a human interferon and a human immunoglobulin Fc fragment, in which the C-terminal of the interferon is linked to the N-terminal of the Fc fragment by a peptide linker moiety. An example of a peptide linker is a peptide comprising primarily a T cell inert sequence, which is immunologically inert. An exemplary peptide linker has the amino acid sequence: GGS GG SGGGG SGGGG S (SEQ ID NO:4). In such a fusion protein, an illustrative Fc moiety is a human $\gamma 4$ chain, which is stable in solution and has little or no complement activating activity. Accordingly, the present invention contemplates a Zmim19 fusion protein that comprises a Zmim19 moiety and a human Fc fragment, wherein the C-terminus of the Zmim19 moiety is attached to the N-terminus of the Fc fragment via a peptide linker, such as a peptide consisting of the amino acid sequence of SEQ ID NO:4. The Zmim19 moiety can be a Zmim19 molecule or a fragment thereof.

In another variation, a Zmim19 fusion protein comprises an IgG sequence, a Zmim19 moiety covalently joined to the aminoterminal end of the IgG sequence, and a signal peptide that is covalently joined to the aminoterminal of the Zmim19 moiety, wherein the IgG sequence consists of the following elements in the following order: a hinge region, a CH₂ domain, and a CH₃ domain. Accordingly, the IgG sequence lacks a CH₁ domain. The Zmim19 moiety displays a Zmim19 activity, as described herein, such as the ability to bind with a Zmim19 antibody. This general approach to producing fusion proteins that comprise both antibody and nonantibody portions has been described by LaRochelle *et al.*, EP 742830 (WO 95/21258).

Fusion proteins comprising a Zmim19 moiety and an Fc moiety can be used, for example, as an *in vitro* assay tool. For example, the presence of a Zmim19 substrate or inhibitor in a biological sample can be detected using a Zmim19-antibody fusion protein, in which the Zmim19 moiety is used to target the substrate or inhibitor, and a macromolecule, such as Protein A or anti-Fc antibody, is used to detect the bound fusion protein-receptor complex. Furthermore, such fusion proteins can be used to identify molecules that interfere with the binding of Zmim19 and a substrate.

Another type of Zmim19 fusion protein is useful for therapy, in which a Zmim19 moiety provides an enzymatic component of an adenosine deaminase targeting composition, or a "Zmim19 targeting composition." Such targeting compositions comprise a Zmim19 moiety and a recognition moiety capable of specific binding with a chosen target, such as a cell surface marker. For example, the recognition moiety can be a ligand that binds with a cell surface receptor. One example of a recognition moiety is transferrin, which targets cells bearing a transferrin receptor. Another example of a recognition moiety is epidermal growth factor, which binds to cells expressing its cognate receptor.

Alternatively, the recognition moiety can be an antibody, or antibody fragment, which binds a specific cell surface marker on the target cells. Such antibodies, or antibody fragments, can bind, for example, with tumor associated antigens or infectious agent antigens.

General methods for preparing targeting compositions that comprise an adenosine deaminase moiety are known to those of skill in the art. Such targeting compositions can be prepared as fusion proteins, or by chemically coupling a adenosine deaminase moiety and a recognition moiety. See, for example, Goldenberg, U.S. patent No. 5,698,178, Raines *et al.*, U.S. patent No. 5,840,296, Rybak *et al.*, U.S. patent No. 5,955,073, Youle *et al.*, *Crit. Rev. Ther. Drug Carrier Syst.* 10:1 (1993), Rybak and Newton, *Exp. Cell. Res.* 253:325 (1999), and Suzuki *et al.*, *Nature Biotechnology* 17:265 (1999).

Moreover, using methods described in the art, hybrid Zmim19 fusion proteins can be constructed using regions or domains of the inventive Zmim19 in combination with those of other adenosine deaminases or heterologous proteins (see, for example, Picard, *Cur. Opin. Biology* 5:511 (1994)). These methods allow the
5 determination of the biological importance of larger domains or regions in a polypeptide of interest. Such hybrids may alter reaction kinetics, binding, constrict or expand the substrate specificity, or alter tissue and cellular localization of a polypeptide, and can be applied to polypeptides of unknown structure. For example Horisberger and DiMarco, *Pharmac. Ther.* 66:507 (1995), describe the construction of fusion protein hybrids
10 comprising different interferon- α subtypes, as well as hybrids comprising interferon- α domains from different species.

Fusion proteins can be prepared by methods known to those skilled in the art by preparing each component of the fusion protein and chemically conjugating the components. Alternatively, a polynucleotide encoding both components of the fusion
15 protein in the proper reading frame can be generated using known techniques and expressed by the methods described herein. General methods for enzymatic and chemical cleavage of fusion proteins are described, for example, by Ausubel (1995) at pages 16-19 to 16-25.

20 **6. Zmim19 Analogs and Zmim19 Inhibitors**

One general class of Zmim19 analogs are variants having an amino acid sequence that is a mutation of the amino acid sequence disclosed herein. Another general class of Zmim19 analogs is provided by anti-idiotypic antibodies, and fragments thereof, as described below. Moreover, recombinant antibodies comprising anti-idiotypic
25 variable domains can be used as analogs (see, for example, Monfardini *et al.*, *Proc. Assoc. Am. Physicians* 108:420 (1996)). Since the variable domains of anti-idiotypic Zmim19 antibodies mimic Zmim19, these domains can provide Zmim19 enzymatic activity. Methods of producing anti-idiotypic catalytic antibodies are known to those of skill in the art (see, for example, Joron *et al.*, *Ann. N Y Acad. Sci.* 672:216 (1992),
30 Friboulet *et al.*, *Appl. Biochem. Biotechnol.* 47:229 (1994), and Avasle *et al.*, *Ann. N Y Acad. Sci.* 864:118 (1998)).

Another approach to identifying Zmim19 analogs is provided by the use of combinatorial libraries. Methods for constructing and screening phage display and other combinatorial libraries are provided, for example, by Kay *et al.*, *Phage Display of Peptides and Proteins* (Academic Press 1996), Verdine, U.S. Patent No. 5,783,384, Kay,
35 *et al.*, U.S. Patent No. 5,747,334, and Kauffman *et al.*, U.S. Patent No. 5,723,323.

One illustrative *in vitro* use of Zmim19 and its analogs is to selectively produce β -(D)-2',3'-dideoxyinosine in high yields from an α , β -anomeric mixture, using the general procedure described by Farina *et al.*, U.S. Patent No. 5,011,774. The product, β -(D)-2',3'-dideoxyinosine, is useful as an antiviral and antibiotic agent. Adenosine deaminase genes are useful as selectable markers or reporter genes in recombinant host cells (see, for example, Pelcher, U.S. Patent No. 5,474,929). Adenosine deaminase is also useful to determine the freshness of fish and mollusks (see, for example, Ohashi *et al.*, U.S. Patent No. 4,650,752). Those of skill in the art can devise other uses for molecules having Zmim19 activity.

The activity of a Zmim19 polypeptide, fragment, or analog can be determined using a standard assay that measures adenosine deaminase activity. For example, Bax *et al.*, *Brit. J. Haematol.* 109:549 (2000), describe an adenosine deaminase assay, which detects the deamination of adenosine to inosine. Briefly, a test sample is incubated with 3.75 mM adenosine in 100 mM phosphate buffer for 30 minutes. The reaction is stopped with 40% trichloroacetic acid, and the protein precipitate is removed by centrifugation for one minute at 12,000xg. The supernatant is then extracted with water-saturated diethyl ether (pH 5) to remove the trichloroacetic acid. The reaction products are separated by HPLC using a Waters trimodular system with a Spherisorb ODS2 column and an isocratic tetrabutylammonium chloride and ammonium acetate buffer. As additional examples, Bota *et al.*, *Clin. Chim. Acta* 290:145 (2000), describe a method for assaying adenosine deaminase using response surface methodology, whereas Wu *et al.*, *Anal. Biochem.* 266:133 (1999), describe a colorimetric detection method, which is used following fractionation of a test sample by isoelectric focusing.

Solution *in vitro* assays can be used to identify a Zmim19 substrate or inhibitor. Solid phase systems can also be used to identify a substrate or inhibitor of a Zmim19 polypeptide. For example, a Zmim19 polypeptide or Zmim19 fusion protein can be immobilized onto the surface of a receptor chip of a commercially available biosensor instrument (BIACORE, Biacore AB; Uppsala, Sweden). The use of this instrument is disclosed, for example, by Karlsson, *Immunol. Methods* 145:229 (1991), and Cunningham and Wells, *J. Mol. Biol.* 234:554 (1993).

In brief, a Zmim19 polypeptide or fusion protein is covalently attached, using amine or sulphydryl chemistry, to dextran fibers that are attached to gold film within a flow cell. A test sample is then passed through the cell. If a Zmim19 substrate or inhibitor is present in the sample, it will bind to the immobilized polypeptide or fusion protein, causing a change in the refractive index of the medium, which is detected as a change in surface plasmon resonance of the gold film. This system allows the determination on- and off-rates, from which binding affinity can be calculated, and

assessment of the stoichiometry of binding, as well as the kinetic effects of Zmim19 mutation. This system can also be used to examine antibody-antigen interactions, and the interactions of other complement/anti-complement pairs.

Accordingly, polypeptides of the present invention are useful as targets for identifying modulators of adenosine deaminase activity. More particularly, Zmim19 polypeptides are useful for screening or identifying new adenosine deaminase inhibitors. Such adenosine deaminase inhibitors can be used, for example, as pharmacological agents to improve coronary and cerebral blood vessel circulation, to prevent or to treat renal diseases, and as anti-inflammatory agents (see, for example, Yamada, U.S. Patent No. 5,705,491).

7. Production of Zmim19 Polypeptides in Cultured Cells

The polypeptides of the present invention, including full-length polypeptides, functional fragments, and fusion proteins, can be produced in recombinant host cells following conventional techniques. To express a *Zmim19* gene, a nucleic acid molecule encoding the polypeptide must be operably linked to regulatory sequences that control transcriptional expression in an expression vector and then, introduced into a host cell. In addition to transcriptional regulatory sequences, such as promoters and enhancers, expression vectors can include translational regulatory sequences and a marker gene, which is suitable for selection of cells that carry the expression vector.

Expression vectors that are suitable for production of a foreign protein in eukaryotic cells typically contain (1) prokaryotic DNA elements coding for a bacterial replication origin and an antibiotic resistance marker to provide for the growth and selection of the expression vector in a bacterial host; (2) eukaryotic DNA elements that control initiation of transcription, such as a promoter; and (3) DNA elements that control the processing of transcripts, such as a transcription termination/polyadenylation sequence. As discussed above, expression vectors can also include nucleotide sequences encoding a secretory sequence that directs the heterologous polypeptide into the secretory pathway of a host cell. For example, a Zmim19 expression vector may comprise a *Zmim19* gene and a secretory sequence derived from a *Zmim19* gene or another secreted gene.

Zmim19 proteins of the present invention may be expressed in mammalian cells. Examples of suitable mammalian host cells include African green monkey kidney cells (Vero; ATCC CRL 1587), human embryonic kidney cells (293-HEK; ATCC CRL 1573), baby hamster kidney cells (BHK-21, BHK-570; ATCC CRL 8544, ATCC CRL 10314), canine kidney cells (MDCK; ATCC CCL 34), Chinese

hamster ovary cells (CHO-K1; ATCC CCL61; CHO DG44 (Chasin *et al.*, *Som. Cell. Molec. Genet.* 12:555, 1986)), rat pituitary cells (GH1; ATCC CCL82), HeLa S3 cells (ATCC CCL2.2), rat hepatoma cells (H-4-II-E; ATCC CRL 1548) SV40-transformed monkey kidney cells (COS-1; ATCC CRL 1650) and murine embryonic cells (NIH-3T3; ATCC CRL 1658).

For a mammalian host, the transcriptional and translational regulatory signals may be derived from viral sources, such as adenovirus, bovine papilloma virus, simian virus, or the like, in which the regulatory signals are associated with a particular gene which has a high level of expression. Suitable transcriptional and translational regulatory sequences also can be obtained from mammalian genes, such as actin, collagen, myosin, and metallothionein genes.

Transcriptional regulatory sequences include a promoter region sufficient to direct the initiation of RNA synthesis. Suitable eukaryotic promoters include the promoter of the mouse *metallothionein I* gene (Hamer *et al.*, *J. Molec. Appl. Genet.* 1:273 (1982)), the *TK* promoter of *Herpes* virus (McKnight, *Cell* 31:355 (1982)), the *SV40* early promoter (Benoist *et al.*, *Nature* 290:304 (1981)), the *Rous* sarcoma virus promoter (Gorman *et al.*, *Proc. Nat'l Acad. Sci. USA* 79:6777 (1982)), the cytomegalovirus promoter (Foecking *et al.*, *Gene* 45:101 (1980)), and the mouse mammary tumor virus promoter (see, generally, Etcheverry, "Expression of Engineered Proteins in Mammalian Cell Culture," in *Protein Engineering: Principles and Practice*, Cleland *et al.* (eds.), pages 163-181 (John Wiley & Sons, Inc. 1996)).

Alternatively, a prokaryotic promoter, such as the bacteriophage T3 RNA polymerase promoter, can be used to control *Znml9* gene expression in mammalian cells if the prokaryotic promoter is regulated by a eukaryotic promoter (Zhou *et al.*, *Mol. Cell. Biol.* 10:4529 (1990), and Kaufman *et al.*, *Nucl. Acids Res.* 19:4485 (1991)).

An expression vector can be introduced into host cells using a variety of standard techniques including calcium phosphate transfection, liposome-mediated transfection, microprojectile-mediated delivery, electroporation, and the like. The transfected cells can be selected and propagated to provide recombinant host cells that comprise the expression vector stably integrated in the host cell genome. Techniques for introducing vectors into eukaryotic cells and techniques for selecting such stable transformants using a dominant selectable marker are described, for example, by Ausubel (1995) and by Murray (ed.), *Gene Transfer and Expression Protocols* (Humana Press 1991).

For example, one suitable selectable marker is a gene that provides resistance to the antibiotic neomycin. In this case, selection is carried out in the presence of a neomycin-type drug, such as G-418 or the like. Selection systems can also be used

to increase the expression level of the gene of interest, a process referred to as "amplification." Amplification is carried out by culturing transfectants in the presence of a low level of the selective agent and then increasing the amount of selective agent to select for cells that produce high levels of the products of the introduced genes. An exemplary amplifiable selectable marker is dihydrofolate reductase, which confers resistance to methotrexate. Other drug resistance genes (e.g., hygromycin resistance, multi-drug resistance, puromycin acetyltransferase) can also be used. Alternatively, markers that introduce an altered phenotype, such as green fluorescent protein, or cell surface proteins (e.g., CD4, CD8, Class I MHC, and placental alkaline phosphatase) may be used to sort transfected cells from untransfected cells by such means as FACS sorting or magnetic bead separation technology.

Znim19 polypeptides can also be produced by cultured cells using a viral delivery system. Exemplary viruses for this purpose include adenovirus, herpesvirus, vaccinia virus and adeno-associated virus (AAV). Adenovirus, a double-stranded DNA virus, is currently the best studied gene transfer vector for delivery of heterologous nucleic acid (for a review, see Becker *et al.*, *Meth. Cell Biol.* 43:161 (1994), and Douglas and Curiel, *Science & Medicine* 4:44 (1997)). Advantages of the adenovirus system include the accommodation of relatively large DNA inserts, the ability to grow to high-titer, the ability to infect a broad range of mammalian cell types, and flexibility that allows use with a large number of available vectors containing different promoters.

By deleting portions of the adenovirus genome, larger inserts (up to 7 kb) of heterologous DNA can be accommodated. These inserts can be incorporated into the viral DNA by direct ligation or by homologous recombination with a co-transfected plasmid. An option is to delete the essential *E1* gene from the viral vector, which results in the inability to replicate unless the *E1* gene is provided by the host cell. For example, adenovirus vector infected human 293 cells (ATCC Nos. CRL-1573, 45504, 45505) can be grown as adherent cells or in suspension culture at relatively high cell density to produce significant amounts of protein (see Garnier *et al.*, *Cytotechnol.* 15:145 (1994)).

Znim19 genes may also be expressed in other higher eukaryotic cells, such as avian, fungal, insect, yeast, or plant cells. The baculovirus system provides an efficient means to introduce cloned Znim19 genes into insect cells. Suitable expression vectors are based upon the *Autographa californica* multiple nuclear polyhedrosis virus (AcMNPV), and contain well-known promoters such as *Drosophila* heat shock protein (hsp) 70 promoter, *Autographa californica* nuclear polyhedrosis virus immediate-early gene promoter (*ie-1*) and the delayed early 39K promoter, baculovirus p10 promoter, and the *Drosophila* metallothionein promoter. A second method of making recombinant baculovirus utilizes a transposon-based system described by Luckow (Luckow, *et al.*, J.

Virol. 67:4566 (1993)). This system, which utilizes transfer vectors, is sold in the BAC-to-BAC kit (Life Technologies, Rockville, MD). This system utilizes a transfer vector, PFASTBAC (Life Technologies) containing a Tn7 transposon to move the DNA encoding the Zmim19 polypeptide into a baculovirus genome maintained in *E. coli* as a large plasmid called a "bacmid." See, Hill-Perkins and Possee, *J. Gen. Virol.* 71:971 (1990), Bonning, *et al.*, *J. Gen. Virol.* 75:1551 (1994), and Chazenbalk, and Rapoport, *J. Biol. Chem.* 270:1543 (1995). In addition, transfer vectors can include an in-frame fusion with DNA encoding an epitope tag at the C- or N-terminus of the expressed Zmim19 polypeptide, for example, a Glu-Glu epitope tag (Grussenmeyer *et al.*, *Proc. Nat'l Acad. Sci.* 82:7952 (1985)). Using a technique known in the art, a transfer vector containing a Zmim19 gene is transformed into *E. coli*, and screened for bacmids which contain an interrupted *lacZ* gene indicative of recombinant baculovirus. The bacmid DNA containing the recombinant baculovirus genome is then isolated using common techniques.

The illustrative PFASTBAC vector can be modified to a considerable degree. For example, the polyhedrin promoter can be removed and substituted with the baculovirus basic protein promoter (also known as *Pcor*, p6.9 or MP promoter) which is expressed earlier in the baculovirus infection, and has been shown to be advantageous for expressing secreted proteins (see, for example, Hill-Perkins and Possee, *J. Gen. Virol.* 71:971 (1990), Bonning, *et al.*, *J. Gen. Virol.* 75:1551 (1994), and Chazenbalk and Rapoport, *J. Biol. Chem.* 270:1543 (1995). In such transfer vector constructs, a short or long version of the basic protein promoter can be used. Moreover, transfer vectors can be constructed which replace the native Zmim19 secretory signal sequences with secretory signal sequences derived from insect proteins. For example, a secretory signal sequence from Ecdysteroid Glucosyltransferase (EGT), honey bee Melittin (Invitrogen Corporation; Carlsbad, CA), or baculovirus gp67 (PharMingen; San Diego, CA) can be used in constructs.

The recombinant virus or bacmid is used to transfect host cells. Suitable insect host cells include cell lines derived from IPLB-Sf-21, a *Spodoptera frugiperda* pupal ovarian cell line, such as Sf9 (ATCC CRL 1711), Sf21AE, and Sf21 (Invitrogen Corporation; San Diego, CA), as well as *Drosophila* Schneider-2 cells, and the HIGH FIVEO cell line (Invitrogen) derived from *Trichoplusia ni* (U.S. Patent No. 5,300,435). Commercially available serum-free media can be used to grow and to maintain the cells. Suitable media are Sf900 II™ (Life Technologies) or ESF 921™ (Expression Systems) for the Sf9 cells; and Ex-cellO405™ (JRH Biosciences, Lenexa, KS) or Express FiveO™ (Life Technologies) for the *T. ni* cells. When recombinant virus is used, the cells are typically grown up from an inoculation density of approximately 2.5×10^5 cells to a

density of $1-2 \times 10^6$ cells at which time a recombinant viral stock is added at a multiplicity of infection (MOI) of 0.1 to 10, more typically near 3.

Established techniques for producing recombinant proteins in baculovirus systems are provided by Bailey *et al.*, "Manipulation of Baculovirus Vectors," in *Methods in Molecular Biology, Volume 7: Gene Transfer and Expression Protocols*, Murray (ed.), pages 147-168 (The Humana Press, Inc. 1991), by Patel *et al.*, "The baculovirus expression system," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover *et al.* (eds.), pages 205-244 (Oxford University Press 1995), by Ausubel (1995) at pages 16-37 to 16-57, by Richardson (ed.), *Baculovirus Expression Protocols* (The Humana Press, Inc. 1995), and by Lucknow, "Insect Cell Expression Technology," in *Protein Engineering: Principles and Practice*, Cleland *et al.* (eds.), pages 183-218 (John Wiley & Sons, Inc. 1996).

Fungal cells, including yeast cells, can also be used to express the genes described herein. Yeast species of particular interest in this regard include *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Pichia methanolica*. Suitable promoters for expression in yeast include promoters from *GAL1* (galactose), *PGK* (phosphoglycerate kinase), *ADH* (alcohol dehydrogenase), *AOX1* (alcohol oxidase), *HIS4* (histidinol dehydrogenase), and the like. Many yeast cloning vectors have been designed and are readily available. These vectors include YIp-based vectors, such as YIp5, YRp vectors, such as YRp17, YE_p vectors such as YE_p13 and YC_p vectors, such as YC_p19. Methods for transforming *S. cerevisiae* cells with exogenous DNA and producing recombinant polypeptides therefrom are disclosed by, for example, Kawasaki, U.S. Patent No. 4,599,311, Kawasaki *et al.*, U.S. Patent No. 4,931,373, Brake, U.S. Patent No. 4,870,008, Welch *et al.*, U.S. Patent No. 5,037,743, and Murray *et al.*, U.S. Patent No. 4,845,075. Transformed cells are selected by phenotype determined by the selectable marker, commonly drug resistance or the ability to grow in the absence of a particular nutrient (*e.g.*, leucine). An illustrative vector system for use in *Saccharomyces cerevisiae* is the *POT1* vector system disclosed by Kawasaki *et al.* (U.S. Patent No. 4,931,373), which allows transformed cells to be selected by growth in glucose-containing media. Additional suitable promoters and terminators for use in yeast include those from glycolytic enzyme genes (see, *e.g.*, Kawasaki, U.S. Patent No. 4,599,311, Kingsman *et al.*, U.S. Patent No. 4,615,974, and Bitter, U.S. Patent No. 4,977,092) and alcohol dehydrogenase genes. See also U.S. Patents Nos. 4,990,446, 5,063,154, 5,139,936, and 4,661,454.

Transformation systems for other yeasts, including *Hansenula polymorpha*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces fragilis*, *Ustilago maydis*, *Pichia pastoris*, *Pichia methanolica*, *Pichia guilliermondii* and

Candida maltosa are known in the art. See, for example, Gleeson *et al.*, *J. Gen. Microbiol.* 132:3459 (1986), and Cregg, U.S. Patent No. 4,882,279. *Aspergillus* cells may be utilized according to the methods of McKnight *et al.*, U.S. Patent No. 4,935,349. Methods for transforming *Acremonium chrysogenum* are disclosed by Sumino *et al.*, U.S. Patent No. 5,162,228. Methods for transforming *Neurospora* are disclosed by Lambowitz, U.S. Patent No. 4,486,533.

For example, the use of *Pichia methanolica* as host for the production of recombinant proteins is disclosed by Raymond, U.S. Patent No. 5,716,808, Raymond, U.S. Patent No. 5,736,383, Raymond *et al.*, *Yeast* 14:11-23 (1998), and in international publication Nos. WO 97/17450, WO 97/17451, WO 98/02536, and WO 98/02565. DNA molecules for use in transforming *P. methanolica* will commonly be prepared as double-stranded, circular plasmids, which can be linearized prior to transformation. For polypeptide production in *P. methanolica*, the promoter and terminator in the plasmid can be that of a *P. methanolica* gene, such as a *P. methanolica* alcohol utilization gene (*AUG1* or *AUG2*). Other useful promoters include those of the dihydroxyacetone synthase (DHAS), formate dehydrogenase (FMD), and catalase (CAT) genes. To facilitate integration of the DNA into the host chromosome, the entire expression segment of the plasmid can be flanked at both ends by host DNA sequences. An illustrative selectable marker for use in *Pichia methanolica* is a *P. methanolica* *ADE2* gene, which encodes phosphoribosyl-5-aminoimidazole carboxylase (AIRC; EC 4.1.1.21), and which allows *ade2* host cells to grow in the absence of adenine. For large-scale, industrial processes where it is desirable to minimize the use of methanol, host cells can be used in which both methanol utilization genes (*AUG1* and *AUG2*) are deleted. For production of secreted proteins, host cells can be deficient in vacuolar protease genes (*PEP4* and *PRB1*). Electroporation is used to facilitate the introduction of a plasmid containing DNA encoding a polypeptide of interest into *P. methanolica* cells. *P. methanolica* cells can be transformed by electroporation using an exponentially decaying, pulsed electric field having a field strength of from 2.5 to 4.5 kV/cm, preferably about 3.75 kV/cm, and a time constant (t) of from 1 to 40 milliseconds, most preferably about 20 milliseconds.

Expression vectors can also be introduced into plant protoplasts, intact plant tissues, or isolated plant cells. Methods for introducing expression vectors into plant tissue include the direct infection or co-cultivation of plant tissue with *Agrobacterium tumefaciens*, microprojectile-mediated delivery, DNA injection, electroporation, and the like. See, for example, Horsch *et al.*, *Science* 227:1229 (1985), Klein *et al.*, *Biotechnology* 10:268 (1992), and Miki *et al.*, "Procedures for Introducing Foreign DNA into Plants," in

Methods in Plant Molecular Biology and Biotechnology, Glick *et al.* (eds.), pages 67-88 (CRC Press, 1993).

Alternatively, *Zmim19* genes can be expressed in prokaryotic host cells. Suitable promoters that can be used to express *Zmim19* polypeptides in a prokaryotic host are well-known to those of skill in the art and include promoters capable of recognizing the T4, T3, Sp6 and T7 polymerases, the P_R and P_L promoters of bacteriophage lambda, the *trp*, *recA*, heat shock, *lacUV5*, *tac*, *lpp-lacSpr*, *phoA*, and *lacZ* promoters of *E. coli*, promoters of *B. subtilis*, the promoters of the bacteriophages of *Bacillus*, *Streptomyces* promoters, the *int* promoter of bacteriophage lambda, the *bla* promoter of pBR322, and the CAT promoter of the chloramphenicol acetyl transferase gene. Prokaryotic promoters have been reviewed by Glick, *J. Ind. Microbiol.* 1:277 (1987), Watson *et al.*, *Molecular Biology of the Gene*, 4th Ed. (Benjamin Cummins 1987), and by Ausubel *et al.* (1995).

Useful prokaryotic hosts include *E. coli* and *Bacillus subtilis*. Suitable strains of *E. coli* include BL21(DE3), BL21(DE3)pLysS, BL21(DE3)pLysE, DH1, DH4I, DH5, DH5I, DH5IF, DH5MCR, DH10B, DH10B/p3, DH11S, C600, HB101, JM101, JM105, JM109, JM110, K38, RR1, Y1088, Y1089, CSH18, ER1451, and ER1647 (see, for example, Brown (ed.), *Molecular Biology Labfax* (Academic Press 1991)). Suitable strains of *Bacillus subtilis* include BR151, YB886, M1119, M1120, and B170 (see, for example, Hardy, "Bacillus Cloning Methods," in *DNA Cloning: A Practical Approach*, Glover (ed.) (IRL Press 1985)).

When expressing a *Zmim19* polypeptide in bacteria such as *E. coli*, the polypeptide may be retained in the cytoplasm, typically as insoluble granules, or may be directed to the periplasmic space by a bacterial secretion sequence. In the former case, the cells are lysed, and the granules are recovered and denatured using, for example, guanidine isothiocyanate or urea. The denatured polypeptide can then be refolded and dimerized by diluting the denaturant, such as by dialysis against a solution of urea and a combination of reduced and oxidized glutathione, followed by dialysis against a buffered saline solution. In the latter case, the polypeptide can be recovered from the periplasmic space in a soluble and functional form by disrupting the cells (by, for example, sonication or osmotic shock) to release the contents of the periplasmic space and recovering the protein, thereby obviating the need for denaturation and refolding.

Methods for expressing proteins in prokaryotic hosts are well-known to those of skill in the art (see, for example, Williams *et al.*, "Expression of foreign proteins in *E. coli* using plasmid vectors and purification of specific polyclonal antibodies," in *DNA Cloning 2: Expression Systems*, 2nd Edition, Glover *et al.* (eds.), page 15 (Oxford University Press 1995), Ward *et al.*, "Genetic Manipulation and Expression of

Antibodies," in *Monoclonal Antibodies: Principles and Applications*, page 137 (Wiley-Liss, Inc. 1995), and Georgiou, "Expression of Proteins in Bacteria," in *Protein Engineering: Principles and Practice*, Cleland *et al.* (eds.), page 101 (John Wiley & Sons, Inc. 1996)).

5 Standard methods for introducing expression vectors into bacterial, yeast, insect, and plant cells are provided, for example, by Ausubel (1995).

 General methods for expressing and recovering foreign protein produced by a mammalian cell system are provided by, for example, Etcheverry, "Expression of Engineered Proteins in Mammalian Cell Culture," in *Protein Engineering: Principles and Practice*, Cleland *et al.* (eds.), pages 163 (Wiley-Liss, Inc. 1996). Standard techniques for
10 recovering protein produced by a bacterial system is provided by, for example, Grishammer *et al.*, "Purification of over-produced proteins from *E. coli* cells," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover *et al.* (eds.), pages 59-92 (Oxford University Press 1995). Established methods for isolating recombinant proteins from a
15 baculovirus system are described by Richardson (ed.), *Baculovirus Expression Protocols* (The Humana Press, Inc. 1995).

 As an alternative, polypeptides of the present invention can be synthesized by exclusive solid phase synthesis, partial solid phase methods, fragment condensation or classical solution synthesis. These synthesis methods are well-known to those of skill
20 in the art (see, for example, Merrifield, *J. Am. Chem. Soc.* 85:2149 (1963), Stewart *et al.*, "Solid Phase Peptide Synthesis" (2nd Edition), (Pierce Chemical Co. 1984), Bayer and Rapp, *Chem. Pept. Prot.* 3:3 (1986), Atherton *et al.*, *Solid Phase Peptide Synthesis: A Practical Approach* (IRL Press 1989), Fields and Colowick, "Solid-Phase Peptide Synthesis," *Methods in Enzymology Volume 289* (Academic Press 1997), and Lloyd-
25 Williams *et al.*, *Chemical Approaches to the Synthesis of Peptides and Proteins* (CRC Press, Inc. 1997)). Variations in total chemical synthesis strategies, such as "native chemical ligation" and "expressed protein ligation" are also standard (see, for example, Dawson *et al.*, *Science* 266:776 (1994), Hackeng *et al.*, *Proc. Nat'l Acad. Sci. USA* 94:7845 (1997), Dawson, *Methods Enzymol.* 287: 34 (1997), Muir *et al.*, *Proc. Nat'l Acad. Sci. USA* 95:6705 (1998), and Severinov and Muir, *J. Biol. Chem.* 273:16205
30 (1998)).

8. Isolation of Zmim19 Polypeptides

 The polypeptides of the present invention can be purified to at least about
35 80% purity, to at least about 90% purity, to at least about 95% purity, or greater than 95% purity with respect to contaminating macromolecules, particularly other proteins

and nucleic acids, and free of infectious and pyrogenic agents. The polypeptides of the present invention may also be purified to a pharmaceutically pure state, which is greater than 99.9% pure. Certain purified polypeptide preparations are substantially free of other polypeptides, particularly other polypeptides of animal origin.

Fractionation and/or conventional purification methods can be used to obtain preparations of Zmim19 purified from natural sources (*e.g.*, uterine tissue), and recombinant Zmim19 polypeptides and fusion Zmim19 polypeptides purified from recombinant host cells. In general, ammonium sulfate precipitation and acid or chaotropic extraction may be used for fractionation of samples. Exemplary purification steps may include hydroxyapatite, size exclusion, FPLC and reverse-phase high performance liquid chromatography. Suitable chromatographic media include derivatized dextrans, agarose, cellulose, polyacrylamide, specialty silicas, and the like. PEI, DEAE, QAE and Q derivatives are preferred. Exemplary chromatographic media include those media derivatized with phenyl, butyl, or octyl groups, such as Phenyl-Sepharose FF (Pharmacia), Toyopearl butyl 650 (Toso Haas, Montgomeryville, PA), Octyl-Sepharose (Pharmacia) and the like; or polyacrylic resins, such as Amberchrom CG 71 (Toso Haas) and the like. Suitable solid supports include glass beads, silica-based resins, cellulosic resins, agarose beads, cross-linked agarose beads, polystyrene beads, cross-linked polyacrylamide resins and the like that are insoluble under the conditions in which they are to be used. These supports may be modified with reactive groups that allow attachment of proteins by amino groups, carboxyl groups, sulfhydryl groups, hydroxyl groups and/or carbohydrate moieties.

Examples of coupling chemistries include cyanogen bromide activation, N-hydroxysuccinimide activation, epoxide activation, sulfhydryl activation, hydrazide activation, and carboxyl and amino derivatives for carbodiimide coupling chemistries. These and other solid media are well known and widely used in the art, and are available from commercial suppliers. Selection of a particular method for polypeptide isolation and purification is a matter of routine design and is determined in part by the properties of the chosen support. See, for example, *Affinity Chromatography: Principles & Methods* (Pharmacia LKB Biotechnology 1988), and Doonan, *Protein Purification Protocols* (The Humana Press 1996).

Additional variations in Zmim19 isolation and purification can be devised by those of skill in the art. For example, Bota *et al.*, *J. Chromatogr. B. Biomed. Sci. Appl.* 737:237 (2000), describe a method for purifying human adenosine deaminase using ion-exchange chromatography, and purine riboside affinity chromatography. In addition, anti-Zmim19 antibodies, obtained as described below, can be used to isolate large quantities of protein by immunoaffinity purification.

The polypeptides of the present invention can also be isolated by exploitation of particular properties. For example, immobilized metal ion adsorption (IMAC) chromatography can be used to purify histidine-rich proteins, including those comprising polyhistidine tags. Briefly, a gel is first charged with divalent metal ions to form a chelate (Sulkowski, *Trends in Biochem.* 3:1 (1985)). Histidine-rich proteins will be adsorbed to this matrix with differing affinities, depending upon the metal ion used, and will be eluted by competitive elution, lowering the pH, or use of strong chelating agents. Other methods of purification include purification of glycosylated proteins by lectin affinity chromatography and ion exchange chromatography (M. Deutscher, (ed.), *Meth. Enzymol.* 182:529 (1990)). Within additional embodiments of the invention, a fusion of the polypeptide of interest and an affinity tag (e.g., maltose-binding protein, an immunoglobulin domain) may be constructed to facilitate purification.

Zmim19 polypeptides or fragments thereof may also be prepared through chemical synthesis, as described above. Zmim19 polypeptides may be monomers or multimers; glycosylated or non-glycosylated; and may or may not include an initial methionine amino acid residue.

The present invention also contemplates chemically modified Zmim19 compositions, in which a polypeptide comprising a Zmim19 moiety (e.g., a Zmim19 polypeptide or fusion protein) is linked with a polymer. Typically, the polymer is water soluble so that the Zmim19 conjugate does not precipitate in an aqueous environment, such as a physiological environment. An example of a suitable polymer is one that has been modified to have a single reactive group, such as an active ester for acylation, or an aldehyde for alkylation. In this way, the degree of polymerization can be controlled. An example of a reactive aldehyde is polyethylene glycol propionaldehyde, or mono-(C₁-C₁₀) alkoxy, or aryloxy derivatives thereof (see, for example, Harris, *et al.*, U.S. Patent No. 5,252,714). The polymer may be branched or unbranched. Moreover, a mixture of polymers can be used to produce Zmim19 conjugates.

Zmim19 conjugates used for therapy can comprise pharmaceutically acceptable water-soluble polymer moieties. Suitable water-soluble polymers include polyethylene glycol (PEG), monomethoxy-PEG, mono-(C₁-C₁₀)alkoxy-PEG, aryloxy-PEG, poly-(N-vinyl pyrrolidone)PEG, tressyl monomethoxy PEG, PEG propionaldehyde, bis-succinimidyl carbonate PEG, propylene glycol homopolymers, a polypropylene oxide/ethylene oxide co-polymer, polyoxyethylated polyols (e.g., glycerol), polyvinyl alcohol, dextran, cellulose, or other carbohydrate-based polymers. Suitable PEG may have a molecular weight from about 600 to about 60,000, including, for example, 5,000, 12,000, 20,000 and 25,000. A Zmim19 conjugate can also comprise a mixture of such

water-soluble polymers. Anti-Zmim19 antibodies or anti-idiotypic antibodies can also be conjugated with a water-soluble polymer.

One example of a Zmim19 conjugate comprises a Zmim19 moiety and a polyalkyl oxide moiety attached to the *N*-terminus of the Zmim19 moiety. PEG is one suitable polyalkyl oxide. As an illustration, Zmim19 can be modified with PEG, a process known as "PEGylation." PEGylation of Zmim19 can be carried out by any of the PEGylation reactions known in the art (see, for example, EP 0 154 316, Delgado *et al.*, *Crit. Rev. Ther. Drug Carrier Syst* 9:249 (1992), Duncan and Spreafico, *Clin. Pharmacokinet.* 27:290 (1994), and Francis *et al.*, *Int J Hematol* 68:1 (1998)). For example, PEGylation can be performed by an acylation reaction or by an alkylation reaction with a reactive polyethylene glycol molecule. In an alternative approach, Zmim19 conjugates are formed by condensing activated PEG, in which a terminal hydroxy or amino group of PEG has been replaced by an activated linker (see, for example, Karasiewicz *et al.*, U.S. Patent No. 5,382,657).

PEGylation by acylation typically requires reacting an active ester derivative of PEG with a Zmim19 polypeptide. An example of an activated PEG ester is PEG esterified to *N*-hydroxysuccinimide. As used herein, the term "acylation" includes the following types of linkages between Zmim19 and a water soluble polymer: amide, carbamate, urethane, and the like. Methods for preparing PEGylated Zmim19 by acylation will typically comprise the steps of (a) reacting a Zmim19 polypeptide with PEG (such as a reactive ester of an aldehyde derivative of PEG) under conditions whereby one or more PEG groups attach to Zmim19, and (b) obtaining the reaction product(s). Generally, the optimal reaction conditions for acylation reactions will be determined based upon known parameters and desired results. For example, the larger the ratio of PEG: Zmim19, the greater the percentage of polyPEGylated Zmim19 product.

The product of PEGylation by acylation is typically a polyPEGylated Zmim19 product, wherein the lysine ϵ -amino groups are PEGylated via an acyl linking group. An example of a connecting linkage is an amide. Typically, the resulting Zmim19 will be at least 95% mono-, di-, or tri-pegylated, although some species with higher degrees of PEGylation may be formed depending upon the reaction conditions. PEGylated species can be separated from unconjugated Zmim19 polypeptides using standard purification methods, such as dialysis, ultrafiltration, ion exchange chromatography, affinity chromatography, and the like.

PEGylation by alkylation generally involves reacting a terminal aldehyde derivative of PEG with Zmim19 in the presence of a reducing agent. PEG groups are preferably attached to the polypeptide via a $-\text{CH}_2\text{-NH}$ group. Derivatization via

reductive alkylation to produce a monoPEGylated product takes advantage of the differential reactivity of different types of primary amino groups available for derivatization. Typically, the reaction is performed at a pH that allows one to take advantage of the pKa differences between the ϵ -amino groups of the lysine residues and the α -amino group of the *N*-terminal residue of the protein. By such selective derivatization, attachment of a water-soluble polymer that contains a reactive group such as an aldehyde, to a protein is controlled. The conjugation with the polymer occurs predominantly at the *N*-terminus of the protein without significant modification of other reactive groups such as the lysine side chain amino groups. The present invention provides a substantially homogenous preparation of Zmim19 monopolymer conjugates.

Reductive alkylation to produce a substantially homogenous population of monopolymer Zmim19 conjugate molecule can comprise the steps of: (a) reacting a Zmim19 polypeptide with a reactive PEG under reductive alkylation conditions at a pH suitable to permit selective modification of the α -amino group at the amino terminus of the Zmim19, and (b) obtaining the reaction product(s). The reducing agent used for reductive alkylation should be stable in aqueous solution and preferably be able to reduce only the Schiff base formed in the initial process of reductive alkylation. Suitable reducing agents include sodium borohydride, sodium cyanoborohydride, dimethylamine borane, trimethylamine borane, and pyridine borane.

For a substantially homogenous population of monopolymer Zmim19 conjugates, the reductive alkylation reaction conditions are those, which permit the selective attachment of the water soluble polymer moiety to the *N*-terminus of Zmim19. Such reaction conditions generally provide for pKa differences between the lysine amino groups and the α -amino group at the *N*-terminus. The pH also affects the ratio of polymer to protein to be used. In general, if the pH is lower, a larger excess of polymer to protein will be desired because the less reactive the *N*-terminal α -group, the more polymer is needed to achieve optimal conditions. If the pH is higher, the polymer:Zmim19 need not be as large because more reactive groups are available. Typically, the pH will fall within the range of 3 to 9, or 3 to 6.

Another factor to consider is the molecular weight of the water-soluble polymer. Generally, the higher the molecular weight of the polymer, the fewer number of polymer molecules which may be attached to the protein. For PEGylation reactions, the typical molecular weight is about 2 kDa to about 100 kDa, about 5 kDa to about 50 kDa, or about 12 kDa to about 25 kDa. The molar ratio of water-soluble polymer to Zmim19 will generally be in the range of 1:1 to 100:1. Typically, the molar ratio of water-soluble polymer to Zmim19 will be 1:1 to 20:1 for polyPEGylation, and 1:1 to 5:1 for monoPEGylation.

General methods for producing conjugates comprising a therapeutic protein and water-soluble polymer moieties are known in the art. See, for example, Karasiewicz *et al.*, U.S. Patent No. 5,382,657, Greenwald *et al.*, U.S. Patent No. 5,738, 846, Nieforth *et al.*, *Clin. Pharmacol. Ther.* 59:636 (1996), Monkarsch *et al.*, *Anal. Biochem.* 247:434 (1997)).

The present invention contemplates compositions comprising a peptide or polypeptide described herein. Such compositions can further comprise a carrier. The carrier can be a conventional organic or inorganic carrier. Examples of carriers include water, buffer solution, alcohol, propylene glycol, macrogol, sesame oil, corn oil, and the like.

Peptides and polypeptides of the present invention comprise at least six, at least nine, or at least 15 contiguous amino acid residues of an amino acid sequence comprising amino acid residues 208 to 294 of SEQ ID NO:2, or an amino acid sequence consisting of SEQ ID NO:2. Within certain embodiments of the invention, the polypeptides comprise 20, 30, 40, 50, 100, or more contiguous residues of these amino acid sequences. Additional polypeptides can comprise at least 20, at least 30, at least 40, or at least 50 contiguous amino acids of amino acid residues 208 to 294 of SEQ ID NO:2, or the amino acid sequence of SEQ ID NO:2. Nucleic acid molecules encoding such polypeptides are useful as polymerase chain reaction primers and probes.

9. Production of Antibodies to Zmim19 Proteins

Antibodies to Zmim19 can be obtained, for example, using as an antigen the product of a Zmim19 expression vector or Zmim19 isolated from a natural source. Particularly useful anti-Zmim19 antibodies "bind specifically" with Zmim19. Antibodies are considered to be specifically binding if the antibodies exhibit at least one of the following two properties: (1) antibodies bind to Zmim19 with a threshold level of binding activity, and (2) antibodies do not significantly cross-react with polypeptides related to Zmim19.

With regard to the first characteristic, antibodies specifically bind if they bind to a Zmim19 polypeptide, peptide or epitope with a binding affinity (K_a) of $10^6 M^{-1}$ or greater, preferably $10^7 M^{-1}$ or greater, more preferably $10^8 M^{-1}$ or greater, and most preferably $10^9 M^{-1}$ or greater. The binding affinity of an antibody can be readily determined by one of ordinary skill in the art, for example, by Scatchard analysis (Scatchard, *Ann. NY Acad. Sci.* 51:660 (1949)). With regard to the second characteristic, antibodies do not significantly cross-react with related polypeptide molecules, for example, if they detect Zmim19, but not known related polypeptides using a standard

Western blot analysis. Examples of known related polypeptides are orthologs and proteins from the same species that are members of a protein family. For example, specifically-binding anti-Zmim19 antibodies bind with Zmim19, but not with known adenosine deaminase proteins.

5 Anti-Zmim19 antibodies can be produced using antigenic Zmim19 epitope-bearing peptides and polypeptides. Antigenic epitope-bearing peptides and polypeptides of the present invention contain a sequence of at least nine, or between 15 to about 30 amino acids contained within SEQ ID NO:2. However, peptides or polypeptides comprising a larger portion of an amino acid sequence of the invention,
10 containing from 30 to 50 amino acids, or any length up to and including the entire amino acid sequence of a polypeptide of the invention, also are useful for inducing antibodies that bind with Zmim19. It is desirable that the amino acid sequence of the epitope-bearing peptide is selected to provide substantial solubility in aqueous solvents (*i.e.*, the sequence includes relatively hydrophilic residues, while hydrophobic residues are preferably avoided). Moreover, amino acid sequences containing proline residues may
15 be also be desirable for antibody production.

 As an illustration, potential antigenic sites in Zmim19 were identified using the Jameson-Wolf method, Jameson and Wolf, *CABIOS* 4:181, (1988), as implemented by the PROTEAN program (version 3.14) of LASERGENE (DNASTAR;
20 Madison, WI). Default parameters were used in this analysis.

 The Jameson-Wolf method predicts potential antigenic determinants by combining six major subroutines for protein structural prediction. Briefly, the Hopp-Woods method, Hopp *et al.*, *Proc. Nat'l Acad. Sci. USA* 78:3824 (1981), was first used to identify amino acid sequences representing areas of greatest local hydrophilicity
25 (parameter: seven residues averaged). In the second step, Emini's method, Emini *et al.*, *J. Virology* 55:836 (1985), was used to calculate surface probabilities (parameter: surface decision threshold (0.6) = 1). Third, the Karplus-Schultz method, Karplus and Schultz, *Naturwissenschaften* 72:212 (1985), was used to predict backbone chain flexibility (parameter: flexibility threshold (0.2) = 1). In the fourth and fifth steps of the analysis,
30 secondary structure predictions were applied to the data using the methods of Chou-Fasman, Chou, "Prediction of Protein Structural Classes from Amino Acid Composition," in *Prediction of Protein Structure and the Principles of Protein Conformation*, Fasman (ed.), pages 549-586 (Plenum Press 1990), and Garnier-Robson, Garnier *et al.*, *J. Mol. Biol.* 120:97 (1978) (Chou-Fasman parameters: conformation table
35 = 64 proteins; α region threshold = 103; β region threshold = 105; Garnier-Robson parameters: α and β decision constants = 0). In the sixth subroutine, flexibility parameters and hydropathy/solvent accessibility factors were combined to determine a

surface contour value, designated as the "antigenic index." Finally, a peak broadening function was applied to the antigenic index, which broadens major surface peaks by adding 20, 40, 60, or 80% of the respective peak value to account for additional free energy derived from the mobility of surface regions relative to interior regions. This calculation was not applied, however, to any major peak that resides in a helical region, since helical regions tend to be less flexible.

The results of this analysis indicated that the following amino acid sequences of SEQ ID NO:2 would provide suitable antigenic molecules: amino acid residues 5 to 15 ("antigenic molecule 1"), amino acid residues 42 to 48 ("antigenic molecule 2"), amino acid residues 56 to 64 ("antigenic molecule 3"), amino acid residues 77 to 83 ("antigenic molecule 4"), amino acid residues 89 to 100 ("antigenic molecule 5"), amino acid residues 104 to 122 ("antigenic molecule 6"), amino acid residues 127 to 137 ("antigenic molecule 7"), amino acid residues 147 to 153 ("antigenic molecule 8"), amino acid residues 180 to 193 ("antigenic molecule 9") amino acid residues 197 to 203 ("antigenic molecule 10"), amino acid residues 212 to 221 ("antigenic molecule 11"), amino acid residues 226 to 232 ("antigenic molecule 12"), amino acid residues 237 to 244 ("antigenic molecule 13"), amino acid residues 263 to 275 ("antigenic molecule 14"), amino acid residues 291 to 297 ("antigenic molecule 15"), amino acid residues 334 to 352 ("antigenic molecule 16"). The present invention contemplates the use of any one of antigenic molecules 1 to 16 to generate antibodies to Zmim19. The present invention also contemplates polypeptides comprising at least one of antigenic molecules 1 to 16.

Another useful fragment of Zmim19 for generating antibodies comprises amino acid residues 208 to 294 of SEQ ID NO:2.

Polyclonal antibodies to recombinant Zmim19 protein or to Zmim19 isolated from natural sources can be prepared using methods well-known to those of skill in the art. Antibodies can also be generated using a Zmim19-glutathione transferase fusion protein, which is similar to a method described by Burrus and McMahon, *Exp. Cell. Res.* 220:363 (1995). General methods for producing polyclonal antibodies are described, for example, by Green *et al.*, "Production of Polyclonal Antisera," in *Immunochemical Protocols* (Manson, ed.), pages 1-5 (Humana Press 1992), and Williams *et al.*, "Expression of foreign proteins in *E. coli* using plasmid vectors and purification of specific polyclonal antibodies," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover *et al.* (eds.), page 15 (Oxford University Press 1995).

The immunogenicity of a Zmim19 polypeptide can be increased through the use of an adjuvant, such as alum (aluminum hydroxide) or Freund's complete or incomplete adjuvant. Polypeptides useful for immunization also include fusion polypeptides, such as fusions of Zmim19 or a portion thereof with an immunoglobulin

polypeptide or with maltose binding protein. The polypeptide immunogen may be a full-length molecule or a portion thereof. If the polypeptide portion is "hapten-like," such portion may be advantageously joined or linked to a macromolecular carrier (such as keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA) or tetanus toxoid) for immunization.

Although polyclonal antibodies are typically raised in animals such as horse, cow, dog, chicken, rat, mouse, rabbit, goat, guinea pig, or sheep, an anti-Zmim19 antibody of the present invention may also be derived from a subhuman primate antibody. General techniques for raising diagnostically and therapeutically useful antibodies in baboons may be found, for example, in Goldenberg *et al.*, international patent publication No. WO 91/11465, and in Losman *et al.*, *Int. J. Cancer* 46:310 (1990).

Alternatively, monoclonal anti-Zmim19 antibodies can be generated. Rodent monoclonal antibodies to specific antigens may be obtained by methods known to those skilled in the art (see, for example, Kohler *et al.*, *Nature* 256:495 (1975), Coligan *et al.* (eds.), *Current Protocols in Immunology*, Vol. 1, pages 2.5.1-2.6.7 (John Wiley & Sons 1991) ["Coligan"], Picklesley *et al.*, "Production of monoclonal antibodies against proteins expressed in *E. coli*," in *DNA Cloning 2: Expression Systems*, 2nd Edition, Glover *et al.* (eds.), page 93 (Oxford University Press 1995)).

Briefly, monoclonal antibodies can be obtained by injecting mice with a composition comprising a *Zmim19* gene product, verifying the presence of antibody production by removing a serum sample, removing the spleen to obtain B-lymphocytes, fusing the B-lymphocytes with myeloma cells to produce hybridomas, cloning the hybridomas, selecting positive clones which produce antibodies to the antigen, culturing the clones that produce antibodies to the antigen, and isolating the antibodies from the hybridoma cultures.

In addition, an anti-Zmim19 antibody of the present invention may be derived from a human monoclonal antibody. Human monoclonal antibodies are obtained from transgenic mice that have been engineered to produce specific human antibodies in response to antigenic challenge. In this technique, elements of the human heavy and light chain locus are introduced into strains of mice derived from embryonic stem cell lines that contain targeted disruptions of the endogenous heavy chain and light chain loci. The transgenic mice can synthesize human antibodies specific for human antigens, and the mice can be used to produce human antibody-secreting hybridomas. Methods for obtaining human antibodies from transgenic mice are described, for example, by Green *et al.*, *Nature Genet.* 7:13 (1994), Lonberg *et al.*, *Nature* 368:856 (1994), and Taylor *et al.*, *Int. Immun.* 6:579 (1994).

Monoclonal antibodies can be isolated and purified from hybridoma cultures by a variety of well-established techniques. Such isolation techniques include affinity chromatography with Protein-A Sepharose, size-exclusion chromatography, and ion-exchange chromatography (see, for example, Coligan at pages 2.7.1-2.7.12 and pages 2.9.1-2.9.3; Baines *et al.*, "Purification of Immunoglobulin G (IgG)," in *Methods in Molecular Biology*, Vol. 10, pages 79-104 (The Humana Press, Inc. 1992)).

For particular uses, it may be desirable to prepare fragments of anti-Zmim19 antibodies. Such antibody fragments can be obtained, for example, by proteolytic hydrolysis of the antibody. Antibody fragments can be obtained by pepsin or papain digestion of whole antibodies by conventional methods. As an illustration, antibody fragments can be produced by enzymatic cleavage of antibodies with pepsin to provide a 5S fragment denoted F(ab')₂. This fragment can be further cleaved using a thiol reducing agent to produce 3.5S Fab' monovalent fragments. Optionally, the cleavage reaction can be performed using a blocking group for the sulfhydryl groups that result from cleavage of disulfide linkages. As an alternative, an enzymatic cleavage using pepsin produces two monovalent Fab fragments and an Fc fragment directly. These methods are described, for example, by Goldenberg, U.S. patent No. 4,331,647, Nisonoff *et al.*, *Arch Biochem. Biophys.* 89:230 (1960), Porter, *Biochem. J.* 73:119 (1959), Edelman *et al.*, in *Methods in Enzymology Vol. 1*, page 422 (Academic Press 1967), and by Coligan at pages 2.8.1-2.8.10 and 2.10.-2.10.4.

Other methods of cleaving antibodies, such as separation of heavy chains to form monovalent light-heavy chain fragments, further cleavage of fragments, or other enzymatic, chemical or genetic techniques may also be used, so long as the fragments bind to the antigen that is recognized by the intact antibody.

For example, Fv fragments comprise an association of V_H and V_L chains. This association can be noncovalent, as described by Inbar *et al.*, *Proc. Nat'l Acad. Sci. USA* 69:2659 (1972). Alternatively, the variable chains can be linked by an intermolecular disulfide bond or cross-linked by chemicals such as glutaraldehyde (see, for example, Sandhu, *Crit. Rev. Biotech.* 12:437 (1992)).

The Fv fragments may comprise V_H and V_L chains which are connected by a peptide linker. These single-chain antigen binding proteins (scFv) are prepared by constructing a structural gene comprising DNA sequences encoding the V_H and V_L domains which are connected by an oligonucleotide. The structural gene is inserted into an expression vector which is subsequently introduced into a host cell, such as *E. coli*. The recombinant host cells synthesize a single polypeptide chain with a linker peptide bridging the two V domains. Methods for producing scFvs are described, for example, by Whitlow *et al.*, *Methods: A Companion to Methods in Enzymology* 2:97 (1991) (also

see, Bird *et al.*, *Science* 242:423 (1988), Ladner *et al.*, U.S. Patent No. 4,946,778, Pack *et al.*, *Bio/Technology* 11:1271 (1993), and Sandhu, *supra*.

As an illustration, a scFV can be obtained by exposing lymphocytes to Zmim19 polypeptide *in vitro*, and selecting antibody display libraries in phage or similar
5 vectors (for instance, through use of immobilized or labeled Zmim19 protein or peptide). Genes encoding polypeptides having potential Zmim19 polypeptide binding domains can be obtained by screening random peptide libraries displayed on phage (phage display) or on bacteria, such as *E. coli*. Nucleotide sequences encoding the polypeptides can be obtained in a number of ways, such as through random mutagenesis and random
10 polynucleotide synthesis. These random peptide display libraries can be used to screen for peptides which interact with a known target which can be a protein or polypeptide, such as a ligand or receptor, a biological or synthetic macromolecule, or organic or inorganic substances. Techniques for creating and screening such random peptide display libraries are known in the art (Ladner *et al.*, U.S. Patent No. 5,223,409, Ladner
15 *et al.*, U.S. Patent No. 4,946,778, Ladner *et al.*, U.S. Patent No. 5,403,484, Ladner *et al.*, U.S. Patent No. 5,571,698, and Kay *et al.*, *Phage Display of Peptides and Proteins* (Academic Press, Inc. 1996)) and random peptide display libraries and kits for screening such libraries are available commercially, for instance from CLONTECH Laboratories, Inc. (Palo Alto, CA), Invitrogen Inc. (San Diego, CA), New England Biolabs, Inc. (Beverly, MA), and Pharmacia LKB Biotechnology Inc. (Piscataway, NJ). Random
20 peptide display libraries can be screened using the Zmim19 sequences disclosed herein to identify proteins which bind to Zmim19.

Another form of an antibody fragment is a peptide coding for a single complementarity-determining region (CDR). CDR peptides ("minimal recognition
25 units") can be obtained by constructing genes encoding the CDR of an antibody of interest. Such genes are prepared, for example, by using the polymerase chain reaction to synthesize the variable region from RNA of antibody-producing cells (see, for example, Larrick *et al.*, *Methods: A Companion to Methods in Enzymology* 2:106 (1991), Courtenay-Luck, "Genetic Manipulation of Monoclonal Antibodies," in *Monoclonal Antibodies: Production, Engineering and Clinical Application*, Ritter *et al.* (eds.), page
30 166 (Cambridge University Press 1995), and Ward *et al.*, "Genetic Manipulation and Expression of Antibodies," in *Monoclonal Antibodies: Principles and Applications*, Birch *et al.*, (eds.), page 137 (Wiley-Liss, Inc. 1995)).

Alternatively, an anti-Zmim19 antibody may be derived from a
35 "humanized" monoclonal antibody. Humanized monoclonal antibodies are produced by transferring mouse complementary determining regions from heavy and light variable chains of the mouse immunoglobulin into a human variable domain. Typical residues of

human antibodies are then substituted in the framework regions of the murine counterparts. The use of antibody components derived from humanized monoclonal antibodies obviates potential problems associated with the immunogenicity of murine constant regions. General techniques for cloning murine immunoglobulin variable domains are described, for example, by Orlandi *et al.*, *Proc. Nat'l Acad. Sci. USA* 86:3833 (1989). Techniques for producing humanized monoclonal antibodies are described, for example, by Jones *et al.*, *Nature* 321:522 (1986), Carter *et al.*, *Proc. Nat'l Acad. Sci. USA* 89:4285 (1992), Sandhu, *Crit. Rev. Biotech.* 12:437 (1992), Singer *et al.*, *J. Immun.* 150:2844 (1993), Sudhir (ed.), *Antibody Engineering Protocols* (Humana Press, Inc. 1995), Kelley, "Engineering Therapeutic Antibodies," in *Protein Engineering: Principles and Practice*, Cleland *et al.* (eds.), pages 399-434 (John Wiley & Sons, Inc. 1996), and by Queen *et al.*, U.S. Patent No. 5,693,762 (1997).

Polyclonal anti-idiotypic antibodies can be prepared by immunizing animals with anti-Zmim19 antibodies or antibody fragments, using standard techniques. See, for example, Green *et al.*, "Production of Polyclonal Antisera," in *Methods In Molecular Biology: Immunochemical Protocols*, Manson (ed.), pages 1-12 (Humana Press 1992). Also, see Coligan at pages 2.4.1-2.4.7. Alternatively, monoclonal anti-idiotypic antibodies can be prepared using anti-Zmim19 antibodies or antibody fragments as immunogens with the techniques, described above. As another alternative, humanized anti-idiotypic antibodies or subhuman primate anti-idiotypic antibodies can be prepared using the above-described techniques. Methods for producing anti-idiotypic antibodies are described, for example, by Irie, U.S. Patent No. 5,208,146, Greene, *et al.*, U.S. Patent No. 5,637,677, and Varthakavi and Minocha, *J. Gen. Virol.* 77:1875 (1996).

Anti-idiotypic Zmim19 antibodies, as well as Zmim19 polypeptides, can be used to identify and to isolate Zmim19 substrates and inhibitors. For example, proteins and peptides of the present invention can be immobilized on a column and used to bind substrate and inhibitor proteins from biological samples that are run over the column (Hermanson *et al.* (eds.), *Immobilized Affinity Ligand Techniques*, pages 195-202 (Academic Press 1992)). Radiolabeled or affinity labeled Zmim19 polypeptides can also be used to identify or to localize Zmim19 substrates and inhibitors in a biological sample (see, for example, Deutscher (ed.), *Methods in Enzymol.*, vol. 182, pages 721-37 (Academic Press 1990); Brunner *et al.*, *Ann. Rev. Biochem.* 62:483 (1993); Fedan *et al.*, *Biochem. Pharmacol.* 33:1167 (1984)).

10. Use of Zmim19 Nucleotide Sequences to Detect Zmim19 Gene Expression and to Examine Zmim19 Gene Structure

Nucleic acid molecules can be used to detect the expression of a Zmim19 gene in a biological sample. Such probe molecules include double-stranded nucleic acid molecules comprising the nucleotide sequence of SEQ ID NO:1, or a portion thereof, as well as single-stranded nucleic acid molecules having the complement of the nucleotide sequence of SEQ ID NO:1, or a portion thereof. As used herein, the term "portion" refers to at least eight nucleotides to at least 20 or more nucleotides. Probe molecules may be DNA, RNA, oligonucleotides, and the like. Certain probes bind with regions of a Zmim19 gene that have a low sequence similarity to comparable regions in other adenosine deaminases. Illustrative biological samples include blood, urine, saliva, tissue biopsy, and autopsy material.

In a basic assay, a single-stranded probe molecule is incubated with RNA, isolated from a biological sample, under conditions of temperature and ionic strength that promote base pairing between the probe and target Zmim19 RNA species. After separating unbound probe from hybridized molecules, the amount of hybrids is detected.

Well-established hybridization methods of RNA detection include northern analysis and dot/slot blot hybridization (see, for example, Ausubel (1995) at pages 4-1 to 4-27, and Wu *et al.* (eds.), "Analysis of Gene Expression at the RNA Level," in *Methods in Gene Biotechnology*, pages 225-239 (CRC Press, Inc. 1997)). Nucleic acid probes can be detectably labeled with radioisotopes such as ³²P or ³⁵S. Alternatively, Zmim19 RNA can be detected with a nonradioactive hybridization method (see, for example, Isaac (ed.), *Protocols for Nucleic Acid Analysis by Nonradioactive Probes* (Humana Press, Inc. 1993)). Typically, nonradioactive detection is achieved by enzymatic conversion of chromogenic or chemiluminescent substrates. Illustrative nonradioactive moieties include biotin, fluorescein, and digoxigenin.

Zmim19 oligonucleotide probes are also useful for *in vivo* diagnosis. As an illustration, ¹⁸F-labeled oligonucleotides can be administered to a subject and visualized by positron emission tomography (Tavitt *et al.*, *Nature Medicine* 4:467 (1998)).

Numerous diagnostic procedures take advantage of the polymerase chain reaction (PCR) to increase sensitivity of detection methods. Standard techniques for performing PCR are well-known (see, generally, Mathew (ed.), *Protocols in Human Molecular Genetics* (Humana Press, Inc. 1991), White (ed.), *PCR Protocols: Current Methods and Applications* (Humana Press, Inc. 1993), Cotter (ed.), *Molecular Diagnosis of Cancer* (Humana Press, Inc. 1996), Hanausek and Walaszek (eds.), *Tumor Marker Protocols* (Humana Press, Inc. 1998), Lo (ed.), *Clinical Applications of PCR* (Humana Press, Inc. 1998), and Meltzer (ed.), *PCR in Bioanalysis* (Humana Press, Inc. 1998)).

Certain PCR primers can be designed to amplify a portion of the *Zmim19* gene that has a low sequence similarity to a comparable region in other adenosine deaminases.

One variation of PCR for diagnostic assays is reverse transcriptase-PCR (RT-PCR). In the RT-PCR technique, RNA is isolated from a biological sample, reverse transcribed to cDNA, and the cDNA is incubated with *Zmim19* primers (see, for example, Wu *et al.* (eds.), "Rapid Isolation of Specific cDNAs or Genes by PCR," in *Methods in Gene Biotechnology*, pages 15-28 (CRC Press, Inc. 1997)). PCR is then performed and the products are analyzed using standard techniques.

As an illustration, RNA is isolated from biological sample using, for example, the guanidinium-thiocyanate cell lysis procedure described above. Alternatively, a solid-phase technique can be used to isolate mRNA from a cell lysate. A reverse transcription reaction can be primed with the isolated RNA using random oligonucleotides, short homopolymers of dT, or *Zmim19* anti-sense oligomers. Oligo-dT primers offer the advantage that various mRNA nucleotide sequences are amplified that can provide control target sequences. *Zmim19* sequences are amplified by the polymerase chain reaction using two flanking oligonucleotide primers that are typically 20 bases in length.

PCR amplification products can be detected using a variety of approaches. For example, PCR products can be fractionated by gel electrophoresis, and visualized by ethidium bromide staining. Alternatively, fractionated PCR products can be transferred to a membrane, hybridized with a detectably-labeled *Zmim19* probe, and examined by autoradiography. Additional alternative approaches include the use of digoxigenin-labeled deoxyribonucleic acid triphosphates to provide chemiluminescence detection, and the C-TRAK colorimetric assay.

Another approach for detection of *Zmim19* expression is cycling probe technology, in which a single-stranded DNA target binds with an excess of DNA-RNA-DNA chimeric probe to form a complex, the RNA portion is cleaved with RNAase H, and the presence of cleaved chimeric probe is detected (see, for example, Beggs *et al.*, *J. Clin. Microbiol.* 34:2985 (1996), Bekkaoui *et al.*, *Biotechniques* 20:240 (1996)). Alternative methods for detection of *Zmim19* sequences can utilize approaches such as nucleic acid sequence-based amplification, cooperative amplification of templates by cross-hybridization, and the ligase chain reaction (see, for example, Marshall *et al.*, U.S. Patent No. 5,686,272 (1997), Dyer *et al.*, *J. Virol. Methods* 60:161 (1996), Ehrlich *et al.*, *Eur. J. Biochem.* 243:358 (1997), and Chadwick *et al.*, *J. Virol. Methods* 70:59 (1998)). Other standard methods are known to those of skill in the art.

Zmim19 probes and primers can also be used to detect and to localize *Zmim19* gene expression in tissue samples. Methods for such *in situ* hybridization are well-known to those of skill in the art (see, for example, Choo (ed.), *In Situ Hybridization Protocols* (Humana Press, Inc. 1994), Wu *et al.* (eds.), "Analysis of Cellular DNA or Abundance of mRNA by Radioactive *In Situ* Hybridization (RISH)," in *Methods in Gene Biotechnology*, pages 259-278 (CRC Press, Inc. 1997), and Wu *et al.* (eds.), "Localization of DNA or Abundance of mRNA by Fluorescence *In Situ* Hybridization (RISH)," in *Methods in Gene Biotechnology*, pages 279-289 (CRC Press, Inc. 1997)). Various additional diagnostic approaches are well-known to those of skill in the art (see, for example, Mathew (ed.), *Protocols in Human Molecular Genetics* (Humana Press, Inc. 1991), Coleman and Tsongalis, *Molecular Diagnostics* (Humana Press, Inc. 1996), and Elles, *Molecular Diagnosis of Genetic Diseases* (Humana Press, Inc., 1996)).

As described above, mutations in *adenosine deaminase* genes are associated with various diseases and disorders, including severe immunodeficiency. Adenosine deaminase deficiency can be caused by mutations that result in the absence of *adenosine deaminase* gene expression, or mutations that result in variant enzymes having a low level of adenosine deaminase activity (see, for example, Valerio *et al.*, *EMBO J.* 5:113 (1986); Berkvens *et al.*, *Nucleic Acids Res.* 15:9365 (1987); Akeson *et al.*, *J. Cell. Biochem.* 39:217 (1989); Hirschhorn *et al.*, *Am. J. Hum. Genet.* 49:878 (1991)). Accordingly, nucleic acid molecules comprising *Zmim19* nucleotide sequences can be used to determine whether a subject's chromosomes contain a mutation in the *Zmim19* gene. Detectable chromosomal aberrations at the *Zmim19* gene locus include, but are not limited to, aneuploidy, gene copy number changes, insertions, deletions, restriction site changes and rearrangements. Of particular interest are genetic alterations that inactivate a *Zmim19* gene.

Aberrations associated with a *Zmim19* locus can be detected using nucleic acid molecules of the present invention by employing molecular genetic techniques, such as restriction fragment length polymorphism analysis, short tandem repeat analysis employing PCR techniques, amplification-refractory mutation system analysis, single-strand conformation polymorphism detection, RNase cleavage methods, denaturing gradient gel electrophoresis, fluorescence-assisted mismatch analysis, and other genetic analysis techniques known in the art (see, for example, Mathew (ed.), *Protocols in Human Molecular Genetics* (Humana Press, Inc. 1991), Marian, *Chest* 108:255 (1995), Coleman and Tsongalis, *Molecular Diagnostics* (Human Press, Inc. 1996), Elles (ed.), *Molecular Diagnosis of Genetic Diseases* (Humana Press, Inc. 1996), Landegren (ed.), *Laboratory Protocols for Mutation Detection* (Oxford University Press 1996), Birren *et al.* (eds.), *Genome Analysis, Vol. 2: Detecting Genes* (Cold Spring Harbor Laboratory

Press 1998), Dracopoli *et al.* (eds.), *Current Protocols in Human Genetics* (John Wiley & Sons 1998), and Richards and Ward, "Molecular Diagnostic Testing," in *Principles of Molecular Medicine*, pages 83-88 (Humana Press, Inc. 1998)).

The protein truncation test is also useful for detecting the inactivation of a gene in which translation-terminating mutations produce only portions of the encoded protein (see, for example, Stoppa-Lyonnet *et al.*, *Blood* 91:3920 (1998)). According to this approach, RNA is isolated from a biological sample, and used to synthesize cDNA. PCR is then used to amplify the *Zmim19* target sequence and to introduce an RNA polymerase promoter, a translation initiation sequence, and an in-frame ATG triplet. PCR products are transcribed using an RNA polymerase, and the transcripts are translated *in vitro* with a T7-coupled reticulocyte lysate system. The translation products are then fractionated by SDS-PAGE to determine the lengths of the translation products. The protein truncation test is described, for example, by Dracopoli *et al.* (eds.), *Current Protocols in Human Genetics*, pages 9.11.1 - 9.11.18 (John Wiley & Sons 1998).

The chromosomal location of the *Zmim19* gene can be determined using radiation hybrid mapping, which is a somatic cell genetic technique developed for constructing high-resolution, contiguous maps of mammalian chromosomes (Cox *et al.*, *Science* 250:245 (1990)). Partial or full knowledge of a gene's sequence allows one to design PCR primers suitable for use with chromosomal radiation hybrid mapping panels. Radiation hybrid mapping panels are commercially available which cover the entire human genome, such as the Stanford G3 RH Panel and the GeneBridge 4 RH Panel (Research Genetics, Inc., Huntsville, AL). These panels enable rapid, PCR-based chromosomal localizations and ordering of genes, sequence-tagged sites, and other nonpolymorphic and polymorphic markers within a region of interest. This includes establishing directly proportional physical distances between newly discovered genes of interest and previously mapped markers.

The present invention also contemplates kits for performing a diagnostic assay for *Zmim19* gene expression or to analyze the *Zmim19* locus of a subject. Such kits comprise nucleic acid probes, such as double-stranded nucleic acid molecules comprising the nucleotide sequence of SEQ ID NO:1, or a portion thereof, as well as single-stranded nucleic acid molecules having the complement of the nucleotide sequence of SEQ ID NO:1, or a portion thereof. Illustrative portions reside within nucleotides 172 to 1236 of SEQ ID NO:1. Probe molecules may be DNA, RNA, oligonucleotides, and the like. Kits may comprise nucleic acid primers for performing PCR.

Such a kit can contain all the necessary elements to perform a nucleic acid diagnostic assay described above. A kit will comprise at least one container comprising a

Zmim19 probe or primer. The kit may also comprise a second container comprising one or more reagents capable of indicating the presence of Zmim19 sequences. Examples of such indicator reagents include detectable labels such as radioactive labels, fluorochromes, chemiluminescent agents, and the like. A kit may also comprise a means
5 for conveying to the user that the Zmim19 probes and primers are used to detect Zmim19 gene expression. For example, written instructions may state that the enclosed nucleic acid molecules can be used to detect either a nucleic acid molecule that encodes Zmim19, or a nucleic acid molecule having a nucleotide sequence that is complementary to a Zmim19-encoding nucleotide sequence, or to analyze chromosomal sequences
10 associated with the Zmim19 locus. The written material can be applied directly to a container, or the written material can be provided in the form of a packaging insert.

11. Use of Anti-Zmim19 Antibodies to Detect Zmim19 Protein

The present invention contemplates the use of anti-Zmim19 antibodies to
15 screen biological samples *in vitro* for the presence of Zmim19. In one type of *in vitro* assay, anti-Zmim19 antibodies are used in liquid phase. For example, the presence of Zmim19 in a biological sample can be tested by mixing the biological sample with a trace amount of labeled Zmim19 and an anti-Zmim19 antibody under conditions that promote binding between Zmim19 and its antibody. Complexes of Zmim19 and anti-Zmim19 in
20 the sample can be separated from the reaction mixture by contacting the complex with an immobilized protein which binds with the antibody, such as an Fc antibody or *Staphylococcus* protein A. The concentration of Zmim19 in the biological sample will be inversely proportional to the amount of labeled Zmim19 bound to the antibody and directly related to the amount of free labeled Zmim19. Illustrative biological samples include
25 blood, urine, saliva, tissue biopsy, and autopsy material.

Alternatively, *in vitro* assays can be performed in which anti-Zmim19 antibody is bound to a solid-phase carrier. For example, antibody can be attached to a polymer, such as aminodextran, in order to link the antibody to an insoluble support such as a polymer-coated bead, a plate or a tube. Other suitable *in vitro* assays will be readily
30 apparent to those of skill in the art.

In another approach, anti-Zmim19 antibodies can be used to detect Zmim19 in tissue sections prepared from a biopsy specimen. Such immunochemical detection can be used to determine the relative abundance of Zmim19 and to determine the distribution of Zmim19 in the examined tissue. General immunochemistry techniques are well
35 established (see, for example, Ponder, "Cell Marking Techniques and Their Application," in *Mammalian Development: A Practical Approach*, Monk (ed.), pages 115-38 (IRL Press

1987), Coligan at pages 5.8.1-5.8.8, Ausubel (1995) at pages 14.6.1 to 14.6.13 (Wiley Interscience 1990), and Manson (ed.), *Methods In Molecular Biology, Vol.10: Immunochemical Protocols* (The Humana Press, Inc. 1992)).

Immunochemical detection can be performed by contacting a biological sample with an anti-Zmim19 antibody, and then contacting the biological sample with a detectably labeled molecule, which binds to the antibody. For example, the detectably labeled molecule can comprise an antibody moiety that binds to anti-Zmim19 antibody. Alternatively, the anti-Zmim19 antibody can be conjugated with avidin/streptavidin (or biotin) and the detectably labeled molecule can comprise biotin (or avidin/streptavidin). Numerous variations of this basic technique are well-known to those of skill in the art.

Alternatively, an anti-Zmim19 antibody can be conjugated with a detectable label to form an anti-Zmim19 immunoconjugate. Suitable detectable labels include, for example, a radioisotope, a fluorescent label, a chemiluminescent label, an enzyme label, a bioluminescent label or colloidal gold. Methods of making and detecting such detectably-labeled immunoconjugates are well-known to those of ordinary skill in the art, and are described in more detail below.

The detectable label can be a radioisotope that is detected by autoradiography. Isotopes that are particularly useful for the purpose of the present invention are ^3H , ^{125}I , ^{131}I , ^{35}S and ^{14}C .

Anti-Zmim19 immunoconjugates can also be labeled with a fluorescent compound. The presence of a fluorescently-labeled antibody is determined by exposing the immunoconjugate to light of the proper wavelength and detecting the resultant fluorescence. Fluorescent labeling compounds include fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine.

Alternatively, anti-Zmim19 immunoconjugates can be detectably labeled by coupling an antibody component to a chemiluminescent compound. The presence of the chemiluminescent-tagged immunoconjugate is determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of chemiluminescent labeling compounds include luminol, isoluminol, an aromatic acridinium ester, an imidazole, an acridinium salt and an oxalate ester.

Similarly, a bioluminescent compound can be used to label anti-Zmim19 immunoconjugates of the present invention. Bioluminescence is a type of chemiluminescence found in biological systems in which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent protein is determined by detecting the presence of luminescence. Bioluminescent compounds that are useful for labeling include luciferin, luciferase and aequorin.

Alternatively, anti-Zmim19 immunoconjugates can be detectably labeled by linking an anti-Zmim19 antibody component to an enzyme. When the anti-Zmim19-enzyme conjugate is incubated in the presence of the appropriate substrate, the enzyme moiety reacts with the substrate to produce a chemical moiety which can be detected, for example, by spectrophotometric, fluorometric or visual means. Examples of enzymes that can be used to detectably label polyspecific immunoconjugates include β -galactosidase, glucose oxidase, peroxidase and alkaline phosphatase.

Those of skill in the art will know of other suitable labels which can be employed in accordance with the present invention. The binding of marker moieties to anti-Zmim19 antibodies can be accomplished using standard techniques known to the art. Typical methodology in this regard is described by Kennedy *et al.*, *Clin. Chim. Acta* 70:1 (1976), Schurs *et al.*, *Clin. Chim. Acta* 81:1 (1977), Shih *et al.*, *Int'l J. Cancer* 46:1101 (1990), Stein *et al.*, *Cancer Res.* 50:1330 (1990), and Coligan, *supra*.

Moreover, the convenience and versatility of immunochemical detection can be enhanced by using anti-Zmim19 antibodies that have been conjugated with avidin, streptavidin, and biotin (see, for example, Wilchek *et al.* (eds.), "Avidin-Biotin Technology," *Methods In Enzymology*, Vol. 184 (Academic Press 1990), and Bayer *et al.*, "Immunochemical Applications of Avidin-Biotin Technology," in *Methods In Molecular Biology*, Vol. 10, Manson (ed.), pages 149-162 (The Humana Press, Inc. 1992).

Methods for performing immunoassays are well-established. See, for example, Cook and Self, "Monoclonal Antibodies in Diagnostic Immunoassays," in *Monoclonal Antibodies: Production, Engineering, and Clinical Application*, Ritter and Ladyman (eds.), pages 180-208, (Cambridge University Press, 1995), Perry, "The Role of Monoclonal Antibodies in the Advancement of Immunoassay Technology," in *Monoclonal Antibodies: Principles and Applications*, Birch and Lennox (eds.), pages 107-120 (Wiley-Liss, Inc. 1995), and Diamandis, *Immunoassay* (Academic Press, Inc. 1996).

In a related approach, biotin- or FITC-labeled Zmim19 can be used to identify cells that bind Zmim19. Such can binding can be detected, for example, using flow cytometry.

The present invention also contemplates kits for performing an immunological diagnostic assay for *Zmim19* gene expression. Such kits comprise at least one container comprising an anti-Zmim19 antibody, or antibody fragment. A kit may also comprise a second container comprising one or more reagents capable of indicating the presence of Zmim19 antibody or antibody fragments. Examples of such indicator reagents include detectable labels such as a radioactive label, a fluorescent label, a chemiluminescent label, an enzyme label, a bioluminescent label, colloidal gold, and the like. A kit may also comprise a means for conveying to the user that Zmim19 antibodies

or antibody fragments are used to detect Zmim19 protein. For example, written instructions may state that the enclosed antibody or antibody fragment can be used to detect Zmim19. The written material can be applied directly to a container, or the written material can be provided in the form of a packaging insert.

12. *Therapeutic Uses of Polypeptides Having Zmim19 Activity*

The present invention includes the use of proteins, polypeptides, and peptides having Zmim19 activity (such as Zmim19 polypeptides, anti-idiotypic anti-Zmim19 antibodies, and Zmim19 fusion proteins) to a subject who lacks an adequate amount of adenosine deaminase, such as a subject with severe combined immunodeficiency, as discussed above. As another illustration, adenosine deaminase can be administered to treat CD4+ T cell lymphopenia in HIV-infected patients (see, for example, Shorr *et al.*, U.S. Patent No. 5,728,560). In contrast, an antagonist of Zmim19 (e.g., an anti-Zmim19 antibody) can be used to prevent or to treat a disorder associated with excessive expression of adenosine deaminase. For example, adenosine deaminase inhibitors can be used to treat various inflammatory conditions, such as systemic inflammatory response syndrome, endotoxin shock, septic shock, peritonitis, adult respiratory distress syndrome, and burns (see, for example, Law, U.S. Patent No. 6,103,702).

The Zmim19 molecules described herein can be administered to any subject in need of treatment, and the present invention contemplates both veterinary and human therapeutic uses. Illustrative subjects include mammalian subjects, such as farm animals, domestic animals, and human patients. Human or murine Zmim19 polypeptides can be used for these applications.

Generally, the dosage of administered polypeptide, protein or peptide will vary depending upon such factors as the subject's age, weight, height, sex, general medical condition and previous medical history. Typically, it is desirable to provide the recipient with a dosage of a molecule having Zmim19 activity, which is in the range of from about 1 pg/kg to 10 mg/kg (amount of agent/body weight of subject), although a lower or higher dosage also may be administered as circumstances dictate.

Administration of a molecule having Zmim19 activity to a subject can be intravenous, intraarterial, intraperitoneal, intramuscular, subcutaneous, intrapleural, intrathecal, by perfusion through a regional catheter, or by direct intralesional injection. When administering therapeutic proteins by injection, the administration may be by continuous infusion or by single or multiple boluses.

Additional routes of administration include oral, dermal, mucosal-membrane, pulmonary, and transcutaneous. Oral delivery is suitable for polyester microspheres, zein microspheres, proteinoid microspheres, polycyanoacrylate microspheres, and lipid-based systems (see, for example, DiBase and Morrel, "Oral Delivery of Microencapsulated Proteins," in *Protein Delivery: Physical Systems*, Sanders and Hendren (eds.), pages 255-288 (Plenum Press 1997)). The feasibility of an intranasal delivery is exemplified by such a mode of insulin administration (see, for example, Hinchcliffe and Illum, *Adv. Drug Deliv. Rev.* 35:199 (1999)). Dry or liquid particles comprising molecules having Zmim19 activity can be prepared and inhaled with the aid of dry-powder dispersers, liquid aerosol generators, or nebulizers (e.g., Pettit and Gombotz, *TIBTECH* 16:343 (1998); Patton *et al.*, *Adv. Drug Deliv. Rev.* 35:235 (1999)). This approach is illustrated by the AERX diabetes management system, which is a hand-held electronic inhaler that delivers aerosolized insulin into the lungs. Studies have shown that proteins as large as 48,000 kDa have been delivered across skin at therapeutic concentrations with the aid of low-frequency ultrasound, which illustrates the feasibility of transcutaneous administration (Mitrageoti *et al.*, *Science* 269:850 (1995)). Transdermal delivery using electroporation provides another means to administer Zmim19 moieties (Potts *et al.*, *Pharm. Biotechnol.* 10:213 (1997)).

A pharmaceutical composition comprising a protein, polypeptide, or peptide having Zmim19 activity can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby the therapeutic proteins are combined in a mixture with a pharmaceutically acceptable carrier. A composition is said to be a "pharmaceutically acceptable carrier" if its administration can be tolerated by a recipient patient. Sterile phosphate-buffered saline is one example of a pharmaceutically acceptable carrier. Other suitable carriers are well-known to those in the art. See, for example, Gennaro (ed.), *Remington's Pharmaceutical Sciences*, 19th Edition (Mack Publishing Company 1995).

For purposes of therapy, molecules having Zmim19 activity and a pharmaceutically acceptable carrier are administered to a patient in a therapeutically effective amount. A combination of a protein, polypeptide, or peptide having Zmim19 activity and a pharmaceutically acceptable carrier is said to be administered in a "therapeutically effective amount" if the amount administered is physiologically significant. An agent is physiologically significant if its presence results in a detectable change in the physiology of a recipient patient. For example, an agent comprising a Zmim19 moiety is physiologically significant if its presence increases adenosine deaminase activity.

A pharmaceutical composition comprising molecules having Zmim¹⁹ activity can be furnished in liquid form, or in solid form. Liquid forms, including liposome-encapsulated formulations, are illustrated by injectable solutions and oral suspensions. Exemplary solid forms include capsules, tablets, and controlled-release forms, such as a miniosmotic pump or an implant. Other dosage forms can be devised by those skilled in the art, as shown, for example, by Ansel and Popovich, *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 5th Edition (Lea & Febiger 1990), Gennaro (ed.), *Remington's Pharmaceutical Sciences*, 19th Edition (Mack Publishing Company 1995), and by Ranade and Hollinger, *Drug Delivery Systems* (CRC Press 1996).

Liposomes provide one means to deliver therapeutic polypeptides to a subject intravenously, intraperitoneally, intrathecally, intramuscularly, subcutaneously, or via oral administration, inhalation, or intranasal administration. Liposomes are microscopic vesicles that consist of one or more lipid bilayers surrounding aqueous compartments (see, generally, Bakker-Woudenberg *et al.*, *Eur. J. Clin. Microbiol. Infect. Dis.* 12 (Suppl. 1):S61 (1993), Kim, *Drugs* 46:618 (1993), and Ranade, "Site-Specific Drug Delivery Using Liposomes as Carriers," in *Drug Delivery Systems*, Ranade and Hollinger (eds.), pages 3-24 (CRC Press 1995)). Liposomes are similar in composition to cellular membranes and as a result, liposomes can be administered safely and are biodegradable. Depending on the method of preparation, liposomes may be unilamellar or multilamellar, and liposomes can vary in size with diameters ranging from 0.02 μm to greater than 10 μm . A variety of agents can be encapsulated in liposomes: hydrophobic agents partition in the bilayers and hydrophilic agents partition within the inner aqueous space(s) (see, for example, Machy *et al.*, *Liposomes In Cell Biology And Pharmacology* (John Libbey 1987), and Ostro *et al.*, *American J. Hosp. Pharm.* 46:1576 (1989)). Moreover, it is possible to control the therapeutic availability of the encapsulated agent by varying liposome size, the number of bilayers, lipid composition, as well as the charge and surface characteristics of the liposomes.

Liposomes can adsorb to virtually any type of cell and then slowly release the encapsulated agent. Alternatively, an absorbed liposome may be endocytosed by cells that are phagocytic. Endocytosis is followed by intralysosomal degradation of liposomal lipids and release of the encapsulated agents (Scherphof *et al.*, *Ann. N.Y. Acad. Sci.* 446:368 (1985)). After intravenous administration, small liposomes (0.1 to 1.0 μm) are typically taken up by cells of the reticuloendothelial system, located principally in the liver and spleen, whereas liposomes larger than 3.0 μm are deposited in the lung. This preferential uptake of smaller liposomes by the cells of the reticuloendothelial system

has been used to deliver chemotherapeutic agents to macrophages and to tumors of the liver.

The reticuloendothelial system can be circumvented by several methods including saturation with large doses of liposome particles, or selective macrophage inactivation by pharmacological means (Claassen *et al.*, *Biochim. Biophys. Acta* 802:428 (1984)). In addition, incorporation of glycolipid- or polyethylene glycol-derivatized phospholipids into liposome membranes has been shown to result in a significantly reduced uptake by the reticuloendothelial system (Allen *et al.*, *Biochim. Biophys. Acta* 1068:133 (1991); Allen *et al.*, *Biochim. Biophys. Acta* 1150:9 (1993)).

Liposomes can also be prepared to target particular cells or organs by varying phospholipid composition or by inserting receptors or ligands into the liposomes. For example, liposomes, prepared with a high content of a nonionic surfactant, have been used to target the liver (Hayakawa *et al.*, Japanese Patent 04-244,018; Kato *et al.*, *Biol. Pharm. Bull.* 16:960 (1993)). These formulations were prepared by mixing soybean phosphatidylcholine, α -tocopherol, and ethoxylated hydrogenated castor oil (HCO-60) in methanol, concentrating the mixture under vacuum, and then reconstituting the mixture with water. A liposomal formulation of dipalmitoylphosphatidylcholine (DPPC) with a soybean-derived sterylglucoside mixture (SG) and cholesterol (Ch) has also been shown to target the liver (Shimizu *et al.*, *Biol. Pharm. Bull.* 20:881 (1997)).

Alternatively, various recognition molecules can be bound to the surface of the liposome, such as antibodies, antibody fragments, ligands, carbohydrates, vitamins, transport proteins, and the like. For example, liposomes can be modified with branched type galactosyllipid derivatives to target asialoglycoprotein (galactose) receptors, which are exclusively expressed on the surface of liver cells (Kato and Sugiyama, *Crit. Rev. Ther. Drug Carrier Syst.* 14:287 (1997); Murahashi *et al.*, *Biol. Pharm. Bull.* 20:259 (1997)). Similarly, Wu *et al.*, *Hepatology* 27:772 (1998), have shown that labeling liposomes with asialofetuin led to a shortened liposome plasma half-life and greatly enhanced uptake of asialofetuin-labeled liposome by hepatocytes. On the other hand, hepatic accumulation of liposomes comprising branched type galactosyllipid derivatives can be inhibited by preinjection of asialofetuin (Murahashi *et al.*, *Biol. Pharm. Bull.* 20:259 (1997)). Polyaconitylated human serum albumin liposomes provide another approach for targeting liposomes to liver cells (Kamps *et al.*, *Proc. Nat'l Acad. Sci. USA* 94:11681 (1997)). Moreover, Geho, *et al.* U.S. Patent No. 4,603,044, describe a hepatocyte-directed liposome vesicle delivery system, which has specificity for hepatobiliary receptors associated with the specialized metabolic cells of the liver.

Zn²⁺ targeting compositions can be encapsulated within liposomes using standard techniques of protein microencapsulation (see, for example, Anderson *et*

al., Infect. Immun. 31:1099 (1981), Anderson *et al.*, *Cancer Res.* 50:1853 (1990), and Cohen *et al.*, *Biochim. Biophys. Acta* 1063:95 (1991), Alving *et al.* "Preparation and Use of Liposomes in Immunological Studies," in *Liposome Technology*, 2nd Edition, Vol. III, Gregoriadis (ed.), page 317 (CRC Press 1993), Wassef *et al.*, *Meth. Enzymol.* 149:124 (1987)). Suitable liposomes can contain a variety of moieties, such as lipid derivatives of poly(ethylene glycol) (Allen *et al.*, *Biochim. Biophys. Acta* 1150:9 (1993)).

In addition to providing a means of delivering Zmim19 targeting compositions comprising a Zmim19 moiety and covalently-bound recognition molecule, liposomes can be used to provide a means to administer a Zmim19 moiety and a recognition molecule that are not covalently bound to each other. For example, liposomes can be produced that comprise a Zmim19 moiety and a recognition molecule, wherein the recognition molecule, but not necessarily the Zmim19 moiety, resides on the surface of the liposome. As an illustration, the present invention includes liposomes comprising a recognition molecule positioned on the liposome surface to effect delivery of an encapsulated Zmim19 moiety.

Zmim19 polypeptides can also be delivered to a subject, as entrapped enzymes within energy-replete resealed carrier erythrocytes. This general approach is described, for example, by Bax *et al.*, *Brit. J. Haematol.* 109:5409 (2000).

Zmim19 pharmaceutical compositions may be supplied as a kit comprising a container that comprises Zmim19. Zmim19 can be provided in the form of an injectable solution for single or multiple doses, or as a sterile powder that will be reconstituted before injection. Such a kit may further comprise written information on indications and usage of the pharmaceutical composition. Moreover, such information may include a statement that the Zmim19 composition is contraindicated in patients with known hypersensitivity to Zmim19.

13. Therapeutic Uses of Zmim19 Nucleotide Sequences

The present invention includes the use of Zmim19 nucleotide sequences to provide Zmim19 to a subject in need of Zmim19, as discussed above. In addition, a therapeutic expression vector can be provided that inhibits *Zmim19* gene expression, such as an anti-sense molecule, a ribozyme, or an external guide sequence molecule.

There are numerous approaches to introduce a *Zmim19* gene to a subject, including the use of recombinant host cells that express Zmim19, delivery of naked nucleic acid encoding Zmim19, use of a cationic lipid carrier with a nucleic acid molecule that encodes Zmim19, and the use of viruses that express Zmim19, such as recombinant retroviruses, recombinant adeno-associated viruses, recombinant

adenoviruses, and recombinant Herpes simplex viruses (see, for example, Mulligan, *Science* 260:926 (1993), Rosenberg *et al.*, *Science* 242:1575 (1988), LaSalle *et al.*, *Science* 259:988 (1993), Wolff *et al.*, *Science* 247:1465 (1990), Breakfield and Deluca, *The New Biologist* 3:203 (1991)). In an *ex vivo* approach, for example, cells are isolated from a subject, transfected with a vector that expresses a *Znim19* gene, and then transplanted into the subject. For example, *Znim19* nucleotide sequences can be introduced into a subject's peripheral blood T lymphocytes, hematopoietic stem cells, and the like.

In order to effect expression of a *Znim19* gene, an expression vector is constructed in which a nucleotide sequence encoding a *Znim19* gene is operably linked to a core promoter, and optionally a regulatory element, to control gene transcription. The general requirements of an expression vector are described above.

Alternatively, a *Znim19* gene can be delivered using recombinant viral vectors, including for example, adenoviral vectors (*e.g.*, Kass-Eisler *et al.*, *Proc. Nat'l Acad. Sci. USA* 90:11498 (1993), Kolls *et al.*, *Proc. Nat'l Acad. Sci. USA* 91:215 (1994), Li *et al.*, *Hum. Gene Ther.* 4:403 (1993), Vincent *et al.*, *Nat. Genet.* 5:130 (1993), and Zabner *et al.*, *Cell* 75:207 (1993)), adenovirus-associated viral vectors (Flotte *et al.*, *Proc. Nat'l Acad. Sci. USA* 90:10613 (1993)), alphaviruses such as Semliki Forest Virus and Sindbis Virus (Hertz and Huang, *J. Vir.* 66:857 (1992), Raju and Huang, *J. Vir.* 65:2501 (1991), and Xiong *et al.*, *Science* 243:1188 (1989)), herpes viral vectors (*e.g.*, U.S. Patent Nos. 4,769,331, 4,859,587, 5,288,641 and 5,328,688), parvovirus vectors (Koering *et al.*, *Hum. Gene Therap.* 5:457 (1994)), pox virus vectors (Ozaki *et al.*, *Biochem. Biophys. Res. Comm.* 193:653 (1993), Panicali and Paoletti, *Proc. Nat'l Acad. Sci. USA* 79:4927 (1982)), pox viruses, such as canary pox virus or vaccinia virus (Fisher-Hoch *et al.*, *Proc. Nat'l Acad. Sci. USA* 86:317 (1989), and Flexner *et al.*, *Ann. N.Y. Acad. Sci.* 569:86 (1989)), and retroviruses (*e.g.*, Baba *et al.*, *J. Neurosurg* 79:729 (1993), Ram *et al.*, *Cancer Res.* 53:83 (1993), Takamiya *et al.*, *J. Neurosci. Res* 33:493 (1992), Vile and Hart, *Cancer Res.* 53:962 (1993), Vile and Hart, *Cancer Res.* 53:3860 (1993), and Anderson *et al.*, U.S. Patent No. 5,399,346). Within various embodiments, either the viral vector itself, or a viral particle which contains the viral vector may be utilized in the methods and compositions described below.

As an illustration of one system, adenovirus, a double-stranded DNA virus, is a well-characterized gene transfer vector for delivery of a heterologous nucleic acid molecule (for a review, see Becker *et al.*, *Meth. Cell Biol.* 43:161 (1994); Douglas and Curiel, *Science & Medicine* 4:44 (1997)). The adenovirus system offers several advantages including: (i) the ability to accommodate relatively large DNA inserts, (ii) the ability to be grown to high-titer, (iii) the ability to infect a broad range of mammalian

cell types, and (iv) the ability to be used with many different promoters including ubiquitous, tissue specific, and regulatable promoters. In addition, adenoviruses can be administered by intravenous injection, because the viruses are stable in the bloodstream.

Using adenovirus vectors where portions of the adenovirus genome are deleted, inserts are incorporated into the viral DNA by direct ligation or by homologous recombination with a co-transfected plasmid. In an exemplary system, the essential E1 gene is deleted from the viral vector, and the virus will not replicate unless the E1 gene is provided by the host cell. When intravenously administered to intact animals, adenovirus primarily targets the liver. Although an adenoviral delivery system with an E1 gene deletion cannot replicate in the host cells, the host's tissue will express and process an encoded heterologous protein. Host cells will also secrete the heterologous protein if the corresponding gene includes a secretory signal sequence. Secreted proteins will enter the circulation from tissue that expresses the heterologous gene (e.g., the highly vascularized liver).

Moreover, adenoviral vectors containing various deletions of viral genes can be used to reduce or eliminate immune responses to the vector. Such adenoviruses are E1-deleted, and in addition, contain deletions of E2A or E4 (Lusky *et al.*, *J. Virol.* 72:2022 (1998); Raper *et al.*, *Human Gene Therapy* 9:671 (1998)). The deletion of E2b has also been reported to reduce immune responses (Amalfitano *et al.*, *J. Virol.* 72:926 (1998)). By deleting the entire adenovirus genome, very large inserts of heterologous DNA can be accommodated. Generation of so called "gutless" adenoviruses, where all viral genes are deleted, are particularly advantageous for insertion of large inserts of heterologous DNA (for a review, see Yeh. and Perricaudet, *FASEB J.* 11:615 (1997)).

High titer stocks of recombinant viruses capable of expressing a therapeutic gene can be obtained from infected mammalian cells using standard methods. For example, recombinant HSV can be prepared in Vero cells, as described by Brandt *et al.*, *J. Gen. Virol.* 72:2043 (1991), Herold *et al.*, *J. Gen. Virol.* 75:1211 (1994), Visalli and Brandt, *Virology* 185:419 (1991), Grau *et al.*, *Invest. Ophthalmol. Vis. Sci.* 30:2474 (1989), Brandt *et al.*, *J. Virol. Meth.* 36:209 (1992), and by Brown and MacLean (eds.), *HSV Virus Protocols* (Humana Press 1997).

Alternatively, an expression vector comprising a *Zm19* gene can be introduced into a subject's cells by lipofection *in vivo* using liposomes. Synthetic cationic lipids can be used to prepare liposomes for *in vivo* transfection of a gene encoding a marker (Felgner *et al.*, *Proc. Nat'l Acad. Sci. USA* 84:7413 (1987); Mackey *et al.*, *Proc. Nat'l Acad. Sci. USA* 85:8027 (1988)). The use of lipofection to introduce exogenous genes into specific organs *in vivo* has certain practical advantages. Liposomes can be used to direct transfection to particular cell types, which is particularly

advantageous in a tissue with cellular heterogeneity, such as the pancreas, liver, kidney, and brain. Lipids may be chemically coupled to other molecules for the purpose of targeting. Targeted peptides (e.g., hormones or neurotransmitters), proteins such as antibodies, or non-peptide molecules can be coupled to liposomes chemically.

5 Electroporation is another alternative mode of administration of a Zmim19 nucleic acid molecules. For example, Aihara and Miyazaki, *Nature Biotechnology* 16:867 (1998), have demonstrated the use of *in vivo* electroporation for gene transfer into muscle.

10 In an alternative approach to gene therapy, a therapeutic gene may encode a Zmim19 anti-sense RNA that inhibits the expression of Zmim19. Methods of preparing anti-sense constructs are known to those in the art. See, for example, Erickson *et al.*, *Dev. Genet.* 14:274 (1993) [transgenic mice], Augustine *et al.*, *Dev. Genet.* 14:500 (1993) [murine whole embryo culture], and Olson and Gibo, *Exp. Cell Res.* 241:134 (1998) [cultured cells]. Suitable sequences for Zmim19 anti-sense molecules can be
15 derived from the nucleotide sequences of Zmim19 disclosed herein.

Alternatively, an expression vector can be constructed in which a regulatory element is operably linked to a nucleotide sequence that encodes a ribozyme. Ribozymes can be designed to express endonuclease activity that is directed to a certain target sequence in a mRNA molecule (see, for example, Draper and Macejak, U.S.
20 Patent No. 5,496,698, McSwiggen, U.S. Patent No. 5,525,468, Chowrira and McSwiggen, U.S. Patent No. 5,631,359, and Robertson and Goldberg, U.S. Patent No. 5,225,337). In the context of the present invention, ribozymes include nucleotide sequences that bind with Zmim19 mRNA.

In another approach, expression vectors can be constructed in which a
25 regulatory element directs the production of RNA transcripts capable of promoting RNase P-mediated cleavage of mRNA molecules that encode a Zmim19 gene. According to this approach, an external guide sequence can be constructed for directing the endogenous ribozyme, RNase P, to a particular species of intracellular mRNA, which is subsequently cleaved by the cellular ribozyme (see, for example, Altman *et al.*, U.S. Patent No. 5,168,053, Yuan *et al.*, *Science* 263:1269 (1994), Pace *et al.*, international publication No. WO 96/18733, George *et al.*, international publication No. WO 96/21731, and Werner *et al.*, international publication No. WO 97/33991). Preferably, the external guide sequence comprises a ten to fifteen nucleotide sequence complementary to Zmim19 mRNA, and a 3'-NCCA nucleotide sequence, wherein N is preferably a purine. The external guide
30 sequence transcripts bind to the targeted mRNA species by the formation of base pairs between the mRNA and the complementary external guide sequences, thus promoting
35

cleavage of mRNA by RNase P at the nucleotide located at the 5'-side of the base-paired region.

In general, the dosage of a composition comprising a therapeutic vector having a Zmim19 nucleotide acid sequence, such as a recombinant virus, will vary depending upon such factors as the subject's age, weight, height, sex, general medical condition and previous medical history. Suitable routes of administration of therapeutic vectors include intravenous injection, intraarterial injection, intraperitoneal injection, intramuscular injection, intratumoral injection, and injection into a cavity that contains a tumor.

A composition comprising viral vectors, non-viral vectors, or a combination of viral and non-viral vectors of the present invention can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby vectors or viruses are combined in a mixture with a pharmaceutically acceptable carrier. As noted above, a composition, such as phosphate-buffered saline is said to be a "pharmaceutically acceptable carrier" if its administration can be tolerated by a recipient subject. Other suitable carriers are well-known to those in the art (see, for example, *Remington's Pharmaceutical Sciences*, 19th Ed. (Mack Publishing Co. 1995), and *Gilman's the Pharmacological Basis of Therapeutics*, 7th Ed. (MacMillan Publishing Co. 1985)).

For purposes of therapy, a therapeutic gene expression vector, or a recombinant virus comprising such a vector, and a pharmaceutically acceptable carrier are administered to a subject in a therapeutically effective amount. A combination of an expression vector (or virus) and a pharmaceutically acceptable carrier is said to be administered in a "therapeutically effective amount" if the amount administered is physiologically significant. An agent is physiologically significant if its presence results in a detectable change in the physiology of a recipient subject, as discussed above.

When the subject treated with a therapeutic gene expression vector or a recombinant virus is a human, then the therapy is preferably somatic cell gene therapy. That is, the preferred treatment of a human with a therapeutic gene expression vector or a recombinant virus does not entail introducing into cells a nucleic acid molecule that can form part of a human germ line and be passed onto successive generations (*i.e.*, human germ line gene therapy).

In addition to the uses described above, polynucleotides and polypeptides of the present invention are useful as educational tools in laboratory practicum kits for courses related to genetics and molecular biology, protein chemistry, and antibody production and analysis. Due to its unique polynucleotide and polypeptide sequences, molecules of Zmim19 can be used as standards or as "unknowns" for testing purposes.

For example, Zmim19 polynucleotides can be used as an aid, such as, for example, to teach a student how to prepare expression constructs for bacterial, viral, or mammalian expression, including fusion constructs, wherein Zmim19 is the gene to be expressed; for determining the restriction endonuclease cleavage sites of the polynucleotides; 5 determining mRNA and DNA localization of Zmim19 polynucleotides in tissues (*i.e.*, by northern and Southern blotting as well as polymerase chain reaction); and for identifying related polynucleotides and polypeptides by nucleic acid hybridization. As an illustration, students will find that *Hae*III digestion of a nucleic acid molecule consisting of the nucleotide sequence of nucleotides 172 to 1236 of SEQ ID NO:1 provides 10 fragments of about 72 base pairs, 456 base pairs, and 537 base pairs, and that *Bam*HI digestion yields fragments of about 86 base pairs, 340 base pairs, and 639 base pairs.

Zmim19 polypeptides can be used as an aid to teach preparation of antibodies; identifying proteins by western blotting; protein purification; determining the weight of expressed Zmim19 polypeptides as a ratio to total protein expressed; 15 identifying peptide cleavage sites; coupling amino and carboxyl terminal tags; amino acid sequence analysis, as well as, but not limited to monitoring biological activities of both the native and tagged protein (*i.e.*, protease inhibition) *in vitro* and *in vivo*. For example, students will find that digestion of unglycosylated Zmim19 with NTCB yields five fragments having approximate molecular weights of 1073, 6432, 21647, 3549, and 20 7630, whereas digestion of unglycosylated Zmim19 with cyanogen bromide yields fragments having approximate molecular weights of 148, 3969, 1905, 2025, 1859, 3539, and 26919.

Zmim19 polypeptides can also be used to teach analytical skills such as mass spectrometry, circular dichroism to determine conformation, especially of the four 25 alpha helices, x-ray crystallography to determine the three-dimensional structure in atomic detail, nuclear magnetic resonance spectroscopy to reveal the structure of proteins in solution. For example, a kit containing the Zmim19 can be given to the student to analyze. Since the amino acid sequence would be known by the instructor, the protein can be given to the student as a test to determine the skills or develop the skills of the 30 student, the instructor would then know whether or not the student has correctly analyzed the polypeptide. Since every polypeptide is unique, the educational utility of Zmim19 would be unique unto itself.

The antibodies which bind specifically to Zmim19 can be used as a teaching aid to instruct students how to prepare affinity chromatography columns to 35 purify Zmim19, cloning and sequencing the polynucleotide that encodes an antibody and thus as a practicum for teaching a student how to design humanized antibodies. The *Zmim19* gene, polypeptide, or antibody would then be packaged by reagent companies

and sold to educational institutions so that the students gain skill in art of molecular biology. Because each gene and protein is unique, each gene and protein creates unique challenges and learning experiences for students in a lab practicum. Such educational kits containing the *Znim19* gene, polypeptide, or antibody are considered within the scope of the present invention.

14. Production of Transgenic Mice

Transgenic mice can be engineered to over-express the *Znim19* gene in all tissues or under the control of a tissue-specific or tissue-preferred regulatory element. These over-producers of *Znim19* can be used to characterize the phenotype that results from over-expression, and the transgenic animals can serve as models for human disease caused by excess *Znim19*. Transgenic mice that over-express *Znim19* also provide model bioreactors for production of *Znim19* in the milk or blood of larger animals. Methods for producing transgenic mice are well-known to those of skill in the art (see, for example, Jacob, "Expression and Knockout of Interferons in Transgenic Mice," in *Overexpression and Knockout of Cytokines in Transgenic Mice*, Jacob (ed.), pages 111-124 (Academic Press, Ltd. 1994), Monastersky and Robl (eds.), *Strategies in Transgenic Animal Science* (ASM Press 1995), and Abbud and Nilson, "Recombinant Protein Expression in Transgenic Mice," in *Gene Expression Systems: Using Nature for the Art of Expression*, Fernandez and Hoeffler (eds.), pages 367-397 (Academic Press, Inc. 1999)).

For example, a method for producing a transgenic mouse that expresses a *Znim19* gene can begin with adult, fertile males (studs) (B6C3f1, 2-8 months of age (Taconic Farms, Germantown, NY)), vasectomized males (duds) (B6D2f1, 2-8 months, (Taconic Farms)), prepubescent fertile females (donors) (B6C3f1, 4-5 weeks, (Taconic Farms)) and adult fertile females (recipients) (B6D2f1, 2-4 months, (Taconic Farms)). The donors are acclimated for one week and then injected with approximately 8 IU/mouse of Pregnant Mare's Serum gonadotrophin (Sigma Chemical Company; St. Louis, MO) I.P., and 46-47 hours later, 8 IU/mouse of human Chorionic Gonadotropin (hCG (Sigma)) I.P. to induce superovulation. Donors are mated with studs subsequent to hormone injections. Ovulation generally occurs within 13 hours of hCG injection. Copulation is confirmed by the presence of a vaginal plug the morning following mating.

Fertilized eggs are collected under a surgical scope. The oviducts are collected and eggs are released into urinalysis slides containing hyaluronidase (Sigma). Eggs are washed once in hyaluronidase, and twice in Whitten's W640 medium (described, for example, by Menino and O'Claray, *Biol. Reprod.* 77:159 (1986), and

Dienhart and Downs, *Zygote* 4:129 (1996)) that has been incubated with 5% CO₂, 5% O₂, and 90% N₂ at 37°C. The eggs are then stored in a 37°C/5% CO₂ incubator until microinjection.

5 Ten to twenty micrograms of plasmid DNA containing a Zmim19 encoding sequence is linearized, gel-purified, and resuspended in 10 mM Tris-HCl (pH 7.4), 0.25 mM EDTA (pH 8.0), at a final concentration of 5-10 nanograms per microliter for microinjection. For example, the Zmim19 encoding sequences can encode a polypeptide comprising the amino acid sequence of SEQ ID NO:2 or the amino acid sequence of the murine polypeptide (SEQ ID NO:6).

10 Plasmid DNA is microinjected into harvested eggs contained in a drop of W640 medium overlaid by warm, CO₂-equilibrated mineral oil. The DNA is drawn into an injection needle (pulled from a 0.75mm ID, 1mm OD borosilicate glass capillary), and injected into individual eggs. Each egg is penetrated with the injection needle, into one or both of the haploid pronuclei.

15 Picoliters of DNA are injected into the pronuclei, and the injection needle withdrawn without coming into contact with the nucleoli. The procedure is repeated until all the eggs are injected. Successfully microinjected eggs are transferred into an organ tissue-culture dish with pre-gassed W640 medium for storage overnight in a 37°C/5% CO₂ incubator.

20 The following day, two-cell embryos are transferred into pseudopregnant recipients. The recipients are identified by the presence of copulation plugs, after copulating with vasectomized duds. Recipients are anesthetized and shaved on the dorsal left side and transferred to a surgical microscope. A small incision is made in the skin and through the muscle wall in the middle of the abdominal area outlined by the ribcage, the saddle, and the hind leg, midway between knee and spleen. The reproductive
25 organs are exteriorized onto a small surgical drape. The fat pad is stretched out over the surgical drape, and a baby serrefine (Roboz, Rockville, MD) is attached to the fat pad and left hanging over the back of the mouse, preventing the organs from sliding back in.

30 With a fine transfer pipette containing mineral oil followed by alternating W640 and air bubbles, 12-17 healthy two-cell embryos from the previous day's injection are transferred into the recipient. The swollen ampulla is located and holding the oviduct between the ampulla and the bursa, a nick in the oviduct is made with a 28 g needle close to the bursa, making sure not to tear the ampulla or the bursa.

35 The pipette is transferred into the nick in the oviduct, and the embryos are blown in, allowing the first air bubble to escape the pipette. The fat pad is gently pushed into the peritoneum, and the reproductive organs allowed to slide in. The peritoneal wall

is closed with one suture and the skin closed with a wound clip. The mice recuperate on a 37°C slide warmer for a minimum of four hours.

The recipients are returned to cages in pairs, and allowed 19-21 days gestation. After birth, 19-21 days postpartum is allowed before weaning. The weanlings are sexed and placed into separate sex cages, and a 0.5 cm biopsy (used for genotyping) is snipped off the tail with clean scissors.

Genomic DNA is prepared from the tail snips using, for example, a QIAGEN DNEASY kit following the manufacturer's instructions. Genomic DNA is analyzed by PCR using primers designed to amplify a *Zmim19* gene or a selectable marker gene that was introduced in the same plasmid. After animals are confirmed to be transgenic, they are back-crossed into an inbred strain by placing a transgenic female with a wild-type male, or a transgenic male with one or two wild-type female(s). As pups are born and weaned, the sexes are separated, and their tails snipped for genotyping.

To check for expression of a transgene in a live animal, a partial hepatectomy is performed. A surgical prep is made of the upper abdomen directly below the zyphoid process. Using sterile technique, a small 1.5-2 cm incision is made below the sternum and the left lateral lobe of the liver exteriorized. Using 4-0 silk, a tie is made around the lower lobe securing it outside the body cavity. An atraumatic clamp is used to hold the tie while a second loop of absorbable Dexon (American Cyanamid; Wayne, N.J.) is placed proximal to the first tie. A distal cut is made from the Dexon tie and approximately 100 mg of the excised liver tissue is placed in a sterile petri dish. The excised liver section is transferred to a 14 ml polypropylene round bottom tube and snap frozen in liquid nitrogen and then stored on dry ice. The surgical site is closed with suture and wound clips, and the animal's cage placed on a 37°C heating pad for 24 hours post operatively. The animal is checked daily post operatively and the wound clips removed 7-10 days after surgery. The expression level of *Zmim19* mRNA is examined for each transgenic mouse using an RNA solution hybridization assay or polymerase chain reaction.

In addition to producing transgenic mice that over-express *Zmim19*, it is useful to engineer transgenic mice with either abnormally low or no expression of the gene. Such transgenic mice provide useful models for diseases associated with a lack of *Zmim19*. As discussed above, *Zmim19* gene expression can be inhibited using antisense genes, ribozyme genes, or external guide sequence genes. To produce transgenic mice that under-express the *Zmim19* gene, such inhibitory sequences are targeted to *Zmim19* mRNA. Methods for producing transgenic mice that have abnormally low expression of a particular gene are known to those in the art (see, for example, Wu *et al.*, "Gene Underexpression in Cultured Cells and Animals by Antisense DNA and RNA

Strategies,” in *Methods in Gene Biotechnology*, pages 205-224 (CRC Press 1997)). For example, Blackburn *et al.*, *J. Biol. Chem.* 273:5093 (1998), describe the production of adenosine deaminase-deficient mice using a two-stage genetic engineering approach.

An alternative approach to producing transgenic mice that have little or no *Zmim19* gene expression is to generate mice having at least one normal *Zmim19* allele replaced by a nonfunctional *Zmim19* gene. One method of designing a nonfunctional *Zmim19* gene is to insert another gene, such as a selectable marker gene, within a nucleic acid molecule that encodes *Zmim19*. Standard methods for producing these so-called “knockout mice” are known to those skilled in the art (see, for example, Jacob, “Expression and Knockout of Interferons in Transgenic Mice,” in *Overexpression and Knockout of Cytokines in Transgenic Mice*, Jacob (ed.), pages 111-124 (Academic Press, Ltd. 1994), and Wu *et al.*, “New Strategies for Gene Knockout,” in *Methods in Gene Biotechnology*, pages 339-365 (CRC Press 1997)).

CLAIMS

What is claimed is:

1. An isolated polypeptide, comprising amino acid residues 208 to 294 of SEQ ID NO:2.
2. The isolated polypeptide of claim 1, wherein the isolated polypeptide comprises the amino acid sequence of SEQ ID NO:2.
3. An isolated nucleic acid molecule that encodes a polypeptide, comprising amino acid residues 208 to 294 of SEQ ID NO:2.
4. The isolated nucleic acid molecule of claim 3, wherein the nucleic acid molecule encodes a polypeptide having the amino acid sequence of SEQ ID NO:2.
5. The isolated nucleic acid molecule of claim 4, wherein the nucleic acid molecule comprises the nucleotide sequence of nucleotides 172 to 1236 of SEQ ID NO:1.
6. The isolated nucleic acid molecule of claim 5, wherein the nucleic acid molecule comprises the nucleotide sequence of SEQ ID NO:1.
7. A vector, comprising the isolated nucleic acid molecule of either claim 3 or claim 4.
8. An expression vector, comprising a nucleic acid molecule that encodes the amino acid sequence of SEQ ID NO:2, a transcription promoter, and a transcription terminator, wherein the promoter is operably linked with the nucleic acid molecule, and wherein the nucleic acid molecule is operably linked with the transcription terminator.
9. A recombinant host cell comprising the expression vector of claim 8, wherein the host cell is selected from the group consisting of bacterium, yeast cell, fungal cell, insect cell, mammalian cell, avian cell, and plant cell.
10. A method of using the expression vector of claim 8 to produce a polypeptide that has the amino acid sequence of SEQ ID NO:2, comprising culturing recombinant host cells that comprise the expression vector and that produce the polypeptide.

11. The method of claim 10, further comprising isolating the polypeptide from the cultured recombinant host cells.

12. An antibody or antibody fragment that specifically binds with a polypeptide that has an amino acid sequence consisting of the amino acid sequence of SEQ ID NO:2.

13. A method of detecting in a biological sample the presence of a nucleic acid molecule that encodes the amino acid sequence of SEQ ID NO:2, comprising:

(a) contacting a nucleic acid probe under hybridizing conditions with either (i) test RNA molecules isolated from the biological sample, or (ii) nucleic acid molecules synthesized from the isolated RNA molecules, wherein the probe consists of a nucleotide sequence comprising a portion of the nucleotide sequence of nucleotides 172 to 1236 of SEQ ID NO:1, or a complement thereof, and

(b) detecting the formation of hybrids of the nucleic acid probe and either the test RNA molecules or the synthesized nucleic acid molecules,

wherein the presence of the hybrids indicates the presence of a nucleic acid molecule that encodes the amino acid sequence of SEQ ID NO:2 in the biological sample.

14. A method of detecting in a biological sample the presence of a polypeptide consisting of the amino acid sequence of SEQ ID NO:2, comprising

(a) contacting the biological sample with an antibody, or an antibody fragment, of claim 12, wherein the contacting is performed under conditions that allow the binding of the antibody or antibody fragment to the biological sample, and

(b) detecting any of the bound antibody or bound antibody fragment.

15. A composition, comprising a carrier and the polypeptide of claim 1.

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
21 February 2002 (21.02.2002)

PCT

(10) International Publication Number
WO 02/014483 A3(51) International Patent Classification⁷: C12N 15/55,
9/78, 5/10, C12Q 1/68, G01N 33/53, A61K 38/17

(21) International Application Number: PCT/US01/25457

(22) International Filing Date: 14 August 2001 (14.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/225,493 15 August 2000 (15.08.2000) US(71) Applicant: ZYMOGENETICS INC. [US/US]; 1201
Eastlake Avenue East, Seattle, WA 98102 (US).(72) Inventors: CONKLIN, Darrell, C.; Apt.#7, 2332 Minor
Avenue E, Seattle, WA 98102 (US). GAO, Zeren: 9502
179th Place NE, #3, Redmond, WA 98052 (US).(74) Agent: JONES, Phillip B, C; ZymoGenetics, Inc., 1201
Eastlake Avenue East, Seattle, WA 98102 (US).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW); Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM); European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR); OAPI patent (BF, BJ, CI,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

- with international search report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(88) Date of publication of the international search report:
24 April 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/014483 A3

(54) Title: HUMAN ADENOSINE DEAMINASE

(57) Abstract: Adenosine deaminase is a purine catabolic enzyme, which catalyzes the deamination of adenosine and 2-deoxyadenosine to inosine and 2-deoxyinosine, respectively. A deficiency of adenosine deaminase causes severe immunodeficiency disease. Current therapies for adenosine deficiency include bone marrow transplantation and enzyme replacement therapy. Zm19 is a new human adenosine deaminase, which has uses in therapy, diagnostics, and industry.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 01/25457

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/55 C12N9/78 C12N5/10 C12Q1/68 G01N33/53
A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N G01N C12Q A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, SEQUENCE SEARCH, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

X	<p>DATABASE EMBL SEQUENCE LIBRARY [Online] 6 September 1996 (1996-09-06) HILLIER L., LENNON G., BECKER M., ET AL. : "zk74d09.r1 Soares_pregnant uterus NbHPU Homo sapiens cDNA clone IMAGE:488561"5' similar to WP:C44B7.10 CE02538 ; , mRNA sequence" retrieved from HTTP://SR56.EBI.AC.UK/SRS6BIN/CGI-BIN/WGET Z7-E+[EMBL-ID:HSA44789] Database accession no. AA044789 XP002224431 abstract</p> <p style="text-align: center;">--- -/--</p>	3-6
---	---	-----

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

10 December 2002

Date of mailing of the international search report

27.12.2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 861 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Holtorf, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/25457

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL SEQUENCE LIBRARY [Online] 1 October 1998 (1998-10-01) "NATIONAL CANCER INSTITUTE, CANCER GENOME ANATOMY PROJECT (CGAP), TUMOR GENE INDEX HTTP://WWW.NCBI.NLM.NIH.GOV/NCICGAP": "qf44c04.x1 Soares testis NHT Homo sapiens cDNA clone IMAGE:1752870 3' similar to SW:ADD_ECOLI P22333 ADENOSINE DEAMINASE ;, mRNA sequence" retrieved from HTTP://SRS6.EBI.AC.UK/SRS6BIN/CGI-BIN/WGET Z?-E+[EMBL-ID:A1150008] Database accession no. A1150008 XP002224432 abstract</p> <p>---</p>	3-6
X	<p>DATABASE EMBL SEQUENCE LIBRARY [Online] 26 December 1999 (1999-12-26) ROWEN, L., ET AL. : "Homo sapiens chromosome 15 clone RP11-355D13 map 15q15, complete sequence" retrieved from HTTP://SRS6.EBI.AC.UK/SRS6BIN/CGI-BIN/WGET Z?-E+[EMBL-ID:AC018924] Database accession no. AC018924 XP002224433 abstract</p> <p>---</p>	3-6
P,X	<p>WO 01 57190 A (CAO YICHENG ;CHEN RUI HONG (US); GOODRICH RYLE (US); HYSEQ INC (US) 9 August 2001 (2001-08-09) see SEQID984, and SEQ ID NO 1968 the whole document</p> <p>---</p>	1,3
E	<p>WO 02 26998 A (INCYTE GENOMICS INC ;TRIBOULEY CATHERINE M (US); KHAN FARRAH A (US) 4 April 2002 (2002-04-04) see SEQID5 the whole document</p> <p>---</p>	1-6
E	<p>WO 02 38743 A (GLAXO GROUP LTD ;BURKHART WILLIAM ALFRED (US); MILLER WAYNE HOWARD) 16 May 2002 (2002-05-16) see SEQID1 the whole document</p> <p>---</p>	1,3
E	<p>WO 02 40715 A (CHALUP MICHAEL S;ALTUS CHRISTINA M ; INCYTE GENOMICS INC (US); LIN) 23 May 2002 (2002-05-23) see SEQID62 the whole document</p> <p>-----</p>	1,3

INTERNATIONAL SEARCH REPORT

International Application No.

Information on patent family members

PCT/US 01/25457

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0157190	A	09-08-2001	US 2002128187 A1	12-09-2002
			AU 2591801 A	31-07-2001
			AU 3128801 A	14-08-2001
			AU 3297101 A	07-08-2001
			AU 3300301 A	07-08-2001
			AU 3329301 A	14-08-2001
			AU 3484701 A	14-08-2001
			AU 3484801 A	14-08-2001
			AU 3486501 A	14-08-2001
			AU 3494401 A	14-08-2001
			AU 3665801 A	14-08-2001
			AU 3666001 A	14-08-2001
			AU 3666301 A	14-08-2001
			AU 3672101 A	14-08-2001
			AU 4314201 A	14-08-2001
			EP 1261735 A2	04-12-2002
			EP 1254265 A1	06-11-2002
			EP 1254266 A1	06-11-2002
			EP 1254268 A1	06-11-2002
			EP 1254269 A1	06-11-2002
			EP 1254270 A1	06-11-2002
			WO 0153326 A1	26-07-2001
			WO 0155334 A2	02-08-2001
			WO 0155335 A2	02-08-2001
			WO 0157255 A1	09-08-2001
			WO 0157260 A1	09-08-2001
			WO 0157175 A2	09-08-2001
			WO 0157261 A1	09-08-2001
			WO 0157262 A1	09-08-2001
			WO 0157187 A2	09-08-2001
			WO 0157265 A1	09-08-2001
			WO 0157188 A2	09-08-2001
			WO 0157266 A1	09-08-2001
			WO 0157267 A1	09-08-2001
			WO 0157190 A2	09-08-2001
			US 2002137044 A1	26-09-2002
			US 2002127199 A1	12-09-2002

WO 0226998	A	04-04-2002	AU 9482301 A	08-04-2002
			WO 0226998 A2	04-04-2002

WO 0238743	A	16-05-2002	AU 2901102 A	21-05-2002
			WO 0238743 A2	16-05-2002

WO 0240715	A	23-05-2002	AU 8696401 A	22-03-2002
			AU 8702201 A	22-03-2002
			AU 8710801 A	27-05-2002
			WO 0240715 A2	23-05-2002
			AU 8696101 A	22-03-2002
			AU 8701901 A	22-03-2002
			AU 8707801 A	27-05-2002
